


PATENT COOPERATION TREATY

PCT

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

(PCT Article 36 and Rule 70)



Applicant's or agent's file reference XY - Equine PCT	FOR FURTHER ACTION See Notification of Transmittal of International Preliminary Examination Report (Form PCT/IPEA/416)	
International application No. PCT/US99/17165	International filing date (day/month/year) 29 JULY 1999	Priority date (day/month/year) 30 JULY 1998
International Patent Classification (IPC) or national classification and IPC IPC(7): A61K 39/00, 39/395 and US Cl.: 424/811		
Applicant XY, INC.		

1. This international preliminary examination report has been prepared by this International Preliminary Examining Authority and is transmitted to the applicant according to Article 36.
2. This REPORT consists of a total of 4 sheets.
☒ This report is also accompanied by ANNEXES, i.e., sheets of the description, claims and/or drawings which have been amended and are the basis for this report and/or sheets containing rectifications made before this Authority. (see Rule 70.16 and Section 607 of the Administrative Instructions under the PCT).
These annexes consist of a total of 31 sheets.

3. This report contains indications relating to the following items:

- I ☒ Basis of the report
- II ☐ Priority
- III ☐ Non-establishment of report with regard to novelty, inventive step or industrial applicability
- IV ☐ Lack of unity of invention
- V ☒ Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement
- VI ☐ Certain documents cited
- VII ☐ Certain defects in the international application
- VIII ☐ Certain observations on the international application

Date of submission of the demand 24 FEBRUARY 2000	Date of completion of this report 03 NOVEMBER 2000
Name and mailing address of the IPEA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230	Authorized office JANET M. KERR Telephone No. (703) 308-0196

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No.

PCT/US99/17165

I. Basis of the report

1. With regard to the elements of the international application:*

☐ the international application as originally filed

☒ the description:
 pages (See Attached) _____, as originally filed
 pages _____, filed with the demand
 pages _____, filed with the letter of _____

☒ the claims:
 pages (See Attached) _____, as originally filed
 pages _____, as amended (together with any statement) under Article 19
 pages _____, filed with the demand
 pages _____, filed with the letter of _____

☒ the drawings:
 pages (See Attached) _____, as originally filed
 pages _____, filed with the demand
 pages _____, filed with the letter of _____

☒ the sequence listing part of the description:
 pages (See Attached) _____, as originally filed
 pages _____, filed with the demand
 pages _____, filed with the letter of _____

2. With regard to the language, all the elements marked above were available or furnished to this Authority in the language in which the international application was filed, unless otherwise indicated under this item. These elements were available or furnished to this Authority in the following language _____ which is:

- ☐ the language of a translation furnished for the purposes of international search (under Rule 23.1(b)).
- ☐ the language of publication of the international application (under Rule 48.3(b)).
- ☐ the language of the translation furnished for the purposes of international preliminary examination (under Rules 55.2 and/or 55.3).

3. With regard to any nucleotide and/or amino acid sequence disclosed in the international application, the international preliminary examination was carried out on the basis of the sequence listing:

- ☐ contained in the international application in printed form.
- ☐ filed together with the international application in computer readable form.
- ☐ furnished subsequently to this Authority in written form.
- ☐ furnished subsequently to this Authority in computer readable form.
- ☐ The statement that the subsequently furnished written sequence listing does not go beyond the disclosure in the international application as filed has been furnished.
- ☐ The statement that the information recorded in computer readable form is identical to the written sequence listing has been furnished.

4. ☒ The amendments have resulted in the cancellation of:

- ☒ the description, pages NONE
- ☒ the claims, Nos. NONE
- ☒ the drawings, sheets/fig. NONE

5. ☐ This report has been drawn as if (some of) the amendments had not been made, since they have been considered to go beyond the disclosure as filed, as indicated in the Supplemental Box (Rule 70.2(c)).**

* Replacement sheets which have been furnished to the receiving Office in response to an invitation under Article 14 are referred to in this report as "originally filed" and are not annexed to this report since they do not contain amendments (Rules 70.16 and 70.17).

**Any replacement sheet containing such amendments must be referred to under item 1 and annexed to this report.

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No.

PCT/US99/17165

V. Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement

1. statement

Novelty (N)

Claims 1-67, 69-135

YES

Claims 68

NO

Inventive Step (IS)

Claims NONE

YES

Claims 1-135

NO

Industrial Applicability (IA)

Claims 1-135

YES

Claims NONE

NO

2. citations and explanations (Rule 70.7)

Claim 68 lacks novelty under PCT Article 33(2) as being anticipated by Pinkel et al.

Pinkel et al. disclose sexed sperm samples. As the sexed sperm samples of Pinkel et al. are not distinguishable over the sexed sperm samples of the claimed invention, the claims are anticipated. It is argued that Pinkel et al. do not teach sexed sperm samples, rather mammalian sperm nucleic. This is not persuasive as Pinkel et al. teach sexing ejaculated spermatozoa (see page 270, under "Sample Preparation and Staining", the second paragraph under "Results", and Figures 3-4).

Claims 1-67 and 69-135 lack an inventive step under PCT Article 33(3) as being obvious over McKinnon et al. in view of Pickett et al. and Roth.

McKinnon et al. disclose all of the claim designated features with the exception of insemination near the tip of the uterine horn ipsilateral and/or contralateral to the preovulatory follicle; inserting the insemination sample at a location selected from within about one-half inch, one inch or two inches from the tip of the uterine horn. However, Pickett et al. disclose that the aim of any breeding establishment should be to have a sufficient number of spermatozoa capable of fertilization present in the mare's reproductive tract at the site of fertilization when ovulation occurs. Thus, it would be obvious to inseminate near the tip of the uterine horn ipsilateral and/or contralateral to the preovulatory follicle. Pickett et al. also disclose that the volume of raw or extended inseminate to be used can range from 0.6 to 26.8 ml. The above references do not disclose the specific flow cytometric parameters recited in the claims to isolate and use sexed semen samples. However, Roth discloses sperm sorting through the use of flow cytometry for gender selection. The sorting flow rate is adjusted to 2300 sperm/sec for a sorting rate of 70 to 110 sperm/sec. PBS (Continued on Supplemental Sheet.)

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No.

PCT/US99/17165

Supplemental Box

(To be used when the space in any of the preceding boxes is not sufficient)

Sheet 10

Continuation of: Boxes I - VIII

I. BASIS OF REPORT:

This report has been drawn on the basis of the description,
page(s) 1-42, as originally filed.
page(s) NONE, filed with the demand.
and additional amendments:
NONE

This report has been drawn on the basis of the claims,
page(s) NONE, as originally filed.
page(s) NONE, as amended under Article 19.
page(s) NONE, filed with the demand.
and additional amendments:
Pages 43-73, filed with the letter of 06 September 2000.

This report has been drawn on the basis of the drawings,
page(s) 1-2, as originally filed.
page(s) NONE, filed with the demand.
and additional amendments:
NONE

This report has been drawn on the basis of the sequence listing part of the description:
page(s) NONE, as originally filed.
pages(s) NONE, filed with the demand.
and additional amendments:
NONE

V. 2. REASONED STATEMENTS - CITATIONS AND EXPLANATIONS (Continued):

supplemented with 0.1% BAS is used as sheath fluid. The semen extender contains egg yolk. 350000 to 400000 sperm cells are sorted after 1.5 h. Sorted semen samples are centrifuged and the sperm pellet is resuspended in 200 ul fertilization medium, i.e., the sperm are concentrated. Roth also disclose inseminating mammals with Y chromosome-bearing spermatozoa or X-bearing spermatozoa resulting in offspring of the appropriate sex. Thus, as the flow cytometric techniques are known in the art to successfully sex sperm, and utilization of the sexed sperm in artificial insemination protocols have successfully resulted in offspring of the appropriate sex, it would have been obvious to utilize flow cytometric techniques in a method of producing equine offspring of the desired sex.

It is argued that the references do not teach the step of inseminating near the tip of the uterine horn, sorting equine sperm, or sheath fluid or skim milk solutions for collecting equine sperm samples, or sorting sperm at high speeds. The arguments are not persuasive as applicant is arguing the references individually. The combined references teach the elements recited in the claims or provide motivation to modify the method of McKinnon to obtain sexed sperm which are not damaged by the sorting method, inseminating a mare with the sexed sperm at the site of fertilization when ovulation occurs, and generating sexed offspring.

Claims 1-135 meet the criteria set out in PCT Article 33(4), because the prior art does not teach or fairly suggest as the invention provides a method for producing equine offspring of a desired sex.

----- NEW CITATIONS -----

NONE

PATENT COOPERATION TREATY

PCT

NOTIFICATION OF ELECTION

(PCT Rule 61.2)

From the INTERNATIONAL BUREAU

To:

Assistant Commissioner for Patents
United States Patent and Trademark
Office
Box PCT
Washington, D.C. 20231
ETATS-UNIS D'AMERIQUE

in its capacity as elected Office

Date of mailing (day/month/year)
26 April 2000 (26.04.00)

International application No.
PCT/US99/17165

Applicant's or agent's file reference
XY - Equine PCT

International filing date (day/month/year)
29 July 1999 (29.07.99)

Priority date (day/month/year)
30 July 1998 (30.07.98)

Applicant

SQUIRES, Edward, L. et al

1. The designated Office is hereby notified of its election made:

☒ in the demand filed with the International Preliminary Examining Authority on:
24 February 2000 (24.02.00)

☐ in a notice effecting later election filed with the International Bureau on:

2. The election ☒ was
☐ was not

made before the expiration of 19 months from the priority date or, where Rule 32 applies, within the time limit under Rule 32.2(b).

The International Bureau of WIPO
34, chemin des Colombettes
1211 Geneva 20, Switzerland

Facsimile No.: (41-22) 740.14.35

Authorized officer

Olivia RANAIVOJAONA

Telephone No.: (41-22) 338.83.38

M.H

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 : A61K 39/00, 39/395		(11) International Publication Number: WO 00/06193	
A1		(43) International Publication Date: 10 February 2000 (10.02.00)	
(21) International Application Number: PCT/US99/17165		80524 (US). McCUE, Patrick, M. [US/US]; 1200 Cliffrose Court, Fort Collins, CO 80525 (US). SEIDEL, George, E. [US/US]; 3101 Arrowhead Road, LaPorte, CO 80535 (US).	
(22) International Filing Date: 29 July 1999 (29.07.99)		(74) Agent: SANTANGELO, Luke; Santangelo Law Offices, P.C., 125 South Howes, 3rd floor, Fort Collins, CO 80521 (US).	
(30) Priority Data: 60/094,720 30 July 1998 (30.07.98) US 60/113,143 18 December 1998 (18.12.98) US		(81) Designated States: AE, AL, AM, AT, AT (Utility model), AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, CZ (Utility model), DE, DE (Utility model), DK, DK (Utility model), EE, EE (Utility model), ES, FI, FI (Utility model), GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (Utility model), SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Applications US 60/094,720 (CON) Filed on 30 July 1998 (30.07.98) US 60/113,143 (CON) Filed on 18 December 1998 (18.12.98)		Published With international search report.	
(71) Applicants (for all designated States except US): XY, INC. [US/US]; 3801 Rampart Road, ARBL Building, Fort Collins, CO 80523-1683 (US). COLORADO STATE UNIVERSITY through COLORADO STATE UNIVERSITY RESEARCH FOUNDATION (CSURF) [US/US]; P.O. Box 483, Fort Collins, CO 80522 (US).			
(72) Inventors; and (75) Inventors/Applicants (for US only): SQUIRES, Edward, L. [US/US]; 5536 North County Road 3, Fort Collins, CO			
(54) Title: EQUINE SYSTEM FOR NON-SURGICAL ARTIFICIAL INSEMINATION			
(57) Abstract Non-surgical artificial insemination is achieved for sexed and unsexed equines in a commercially practical manner and with dosages of insemination sperm which were not previously thought to be practical for commercial implementation. Practical and field usable techniques for insemination are presented as well as techniques which offer success ratios at level comparable to the existing routine, high dosage unsexed artificial insemination techniques in equines. Improved insemination and sorting systems particularly adapted to use for sex-selected sperm are disclosed.			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon	KR	Republic of Korea	PL	Poland		
CN	China	KZ	Kazakhstan	PT	Portugal		
CU	Cuba	LC	Saint Lucia	RO	Romania		
CZ	Czech Republic	LI	Liechtenstein	RU	Russian Federation		
DE	Germany	LK	Sri Lanka	SD	Sudan		
DK	Denmark	LR	Liberia	SE	Sweden		
EE	Estonia			SG	Singapore		

EQUINE SYSTEM FOR NON-SURGICAL ARTIFICIAL INSEMINATION

I. TECHNICAL FIELD

5 This invention relates generally to the field of artificial insemination of equines. It also involves equine artificial insemination when there has been sex selection of the sperm to produce an equine offspring of the desired sex. It is especially relevant to situations where non-surgical equine artificial insemination is desired and also where low dose equine artificial insemination is of practical importance.

II. BACKGROUND

10 Artificial insemination of equine mares has been of importance for many years. Often this has been accomplished surgically. In routine instances where lower dosages of sperm have not been required, it has been accomplished without surgery by artificial insemination, however this has used relatively high numbers of sperm. For routine artificial insemination of the mare $250-500 \times 10^6$ progressively motile sperm (pms) inseminated every other day to a
15 mare in estrus, is usually recommended to achieve maximum fertility. Unfortunately, when inseminating mares with semen from a highly fertile stallion, fertility has decreased as the number of motile sperm has been reduced. Under ideal conditions, a mare has been successfully inseminated with as few as 100×10^6 pms without reducing fertility, however even this number poses challenges. Insemination with low numbers of sperm is necessary
20 when using sorted sexed semen and frozen thawed semen of limited quantity.

Preselecting the sex of offspring in an equine animal is, of course, of interest. Sex preselection following artificial insemination (AI) with low numbers of separate, enriched populations of X- and Y- chromosome bearing sperm that have been separated on the basis of DNA content is currently possible in other species, however, equine species have in some
25 regards proven more difficult. While birth of progeny of the desired sex following intrauterine

insemination of cattle and sheep has validated the sexing technology, until the present invention, it has not been practically applied to equines.

To achieve sex preselection involves separating the X- from the Y- chromosome bearing sperm followed by use for artificial insemination (AI) or for *in vitro* fertilization (IVF) and subsequent embryo transfer. Current high speed flow cytometry enables researchers to sort > 1000 live-sexed sperm/second. Sorting alone, however, is not enough. In order to make semen sexing a practical technique for a commercial equine AI program, a lesser number of motile sperm is required for an insemination dose. In the mare when using fresh semen for AI, a typical insemination dose would contain between 250 - 500 x 10⁶ motile sperm. At the current sorting rate of ~1000 live sperm/second, it would take almost six days to sort one insemination dose! Therefore, a lesser number of motile sperm is required to practically achieve reasonable fertility. Once that is achieved, enhanced fertility with non-surgical insemination of mares with sexed semen is also viewed as practically necessary.

As mentioned, sex preselection involves the use of DNA content and separation of sperm into X- and Y-chromosome bearing populations. Using current high speed flow cytometry enables researchers to sort up to 1000 live sperm/sec of the desired sex with 90% accuracy, which provides adequate numbers of sperm in many species other than equines in a reasonable amount of time. For example, this new technology for sperm sexing has made it a practical technique for artificial insemination in cattle. Since it is practical to sort only a low number of spermatozoa and still maintain sperm viability, one aspect of this invention addresses enhanced pregnancy rates following insemination of 25 x 10⁶ non sorted, progressively motile spermatozoa (pms). This is about one-tenth of what was previously considered optimal for routine operations. Other aspects address even lower numbers. These aspects can have significant economic consequences when one considers its application to celebrated trophy animals such as horses and the like.

As mentioned, one of the fundamental challenges that efforts at sorting X and Y equine sperm has faced is the large numbers of sperm involved. In natural insemination equine sperm are produced by nearly the billion; in artificial insemination less, but still

significantly large numbers of equine sperm are usually used. For instance, equine artificial insemination techniques routinely use two hundred and fifty million to five hundred million sperm. Thus a significant number of sperm have been presumed necessary in an equine artificial insemination environment.

5 As the invention relates to sex selected artificial insemination, many methods have been attempted to achieve the separation of X- and Y-chromosome bearing sperm in other animals. None of these, however, have dealt with aspects peculiar to or specific sorting of equine sperm cells. General sorting methods have ranged from magnetic techniques such as appears disclosed in U.S. Patent No. 4276139 to columnar techniques as appears disclosed in
10 U.S. Patent No. 5514537 to gravimetric techniques as discussed in U.S. Patents No. 3894529, reissue Patent No. 32350, U.S. Patents No. 4092229, 4067965, and 4155831. Electrical properties have also been attempted as shown in U.S. 4083957 as well as a combination of electrical and gravimetric properties as discussed in U.S. Patents No. 4225405, 4698142, and 4749458. Motility efforts have also been attempted as shown in U.S. Patents No. 4009260
15 and 4339434. Chemical techniques such as those shown in U.S. Patents No. 4511661 and 4999283 (involving monoclonal antibodies) and U.S. Patents No. 5021244, 5346990, 5439362, and 5660997 (involving membrane proteins), and U.S. Patents No. 3687803, 4191749, 4448767, and 4680258 (involving antibodies) as well as the addition of serum components as shown in U.S. Patent No. 4085205. While each of these techniques has been
20 presented as if to be highly efficient, in fact at present none of those techniques yield the desired level of sex preselection and none have shown success at the artificial insemination level with equine sperm. Regardless of the separation technique eventually used, however, the competing combinations of the high numbers of equine sperm naturally present and the approach of separating X- and Y- chromosome bearing sperm has made it desirable to
25 develop an ability to achieve equine insemination with lower numbers of sperm.

The quantitative technique used to achieve the separation of X-and Y- chromosome bearing sperm for artificial insemination (of any species) has been that involving the technique of flow cytometry. This technique appeared possible as a result of advances and discoveries involving the differential dye absorption of X-and Y- chromosome bearing sperm.

This was discussed early in U.S. Patent No. 4362246 and significantly expanded upon through the techniques disclosed by Lawrence Johnson in U.S. Patent No. 5135759. The Johnson technique of utilizing flow cytometry to separate X- and Y- chromosome bearing sperm has been so significant an advancement that it has for the first time made the commercial separation of such sperm feasible. Further, separation has been significantly enhanced through the utilization of high speed flow cytometers such as the MoFlo® flow cytometer produced by Cytomation, Inc. and discussed in a variety of other patents including US Patent Nos. 5150313, 5602039, 5602349, and 5643796 as well as international PCT patent publication WO 96/12171. While the utilization of Cytomation's MoFlo® cytometers has permitted great increases in speed, and while these speed increases are particularly relevant given the high number of equine sperm often used, certain problems have still remained. In spite of the almost ten-fold advances in speed possible by the MoFlo® flow cytometer, shorter and shorter sorting times have been desired for several reasons. First, it has been discovered that as a practical matter, the equine sperm are time-critical cells. They lose their effectiveness the longer they remain unused. Second, the collection, sorting, and insemination timings has made speed an item of high commercial importance. Thus, the time critical nature of the equine sperm cells and of the process has made speed an essential element in achieving high efficacy and success rates in artificial insemination.

In spite of some successes in sorting and then artificially inseminating animals of other species, the effort with equines has proven particularly elusive. As relevant to the present invention, equine applications may have been particularly challenging either because the equine conception process and/or the equine sperms cells themselves are more delicate than those of other species -- especially bovines. For this reason, it may even be that those skilled in the art have not viewed techniques or systems developed for other species as applicable to equines. In some instances almost identical procedures from a non-equine species do not provide the same type of result for equines. This may have fostered separation in the research efforts and in the techniques and substances developed.

Other problems also exist ranging from the practical to the theoretical. On the practical side, it has been desired to achieve equine artificial insemination in a manner that

can be done in the field rather than a laboratory environment. Thus, for commercial production and success in the field, improvements which might only represent an increase in efficiency or practicality may still be significant. Related to the practical aspect, is the aspect of the delicateness and sensitivity of the entire process. In this regard, it has been desired to simplify the process and make it as procedurally robust as possible so that operator error or skill can play an ever decreasing role. This goal has also combined to make insemination with lower dosages even more desirable.

In addition to the delicateness of the process, it has always been known that the sperm in general are extremely delicate cells. While this factor at first glance seems like it might be considered easily understood, in fact, the full extent of the cells' sensitivities have not yet been fully explored. Furthermore equine sperm appear particularly sensitive. In contrast to bovine sperm, they are in many ways more delicate from the perspective of successful artificial insemination. Different sensitivities arise and thus there has to some degree been a perception that the systems, techniques, and substances used in other animals (such as bovines) may not always be adaptable to equines. This has in fact proven to be true.

In the context of flow cytometry in general, most sorted cells or particles have often been physically able to withstand a variety of abuses. This is not the case for equine sperm cells. In fact, as the present invention discloses, the processing through normal flow cytometer techniques may, in fact, be unacceptable for cytometric sorting of equine sperm cells in certain applications. The sensitivities range from dilution problems and the flow cytometer's inherent need to isolate and distinguish each cell individually as well as the pressure and other stresses which typical flow cytometry has (prior to the present invention) imposed upon the equine cells it was sorting. This may also represent a unique factor for equine sperm cells because it appears that even though the equine sperm cell may appear to pass through the flow cytometer and be sorted with no visually discernable side-effects, in fact, the cells themselves may have been stressed to the point that they perform less than optimally in the insemination process. Thus, an interplay of factors seems involved and has raised unusual problems from the perspective of equine sperm cell sorting and ultimate use for equine artificial insemination.

Another problem which has remained -- in spite of the great advances achieved through the Johnson patent and related technology -- is the fact that prior to the present invention it has been extremely difficult to achieve lower dosage insemination with sexed equine sperm, regardless of the separation technology used. While historically, some achievement of low dose insemination has occurred, it has appeared to be more in a theoretical or laboratory environment rather than in environments which are likely to be experienced in or applicable to a commercial application. It has also occurred through surgical techniques. In this regard, the desire has not been merely to achieve low dose insemination but even to achieve non-surgical insemination in a field environment. To achieve low dose insemination with pregnancy success rates which are comparable to existing unsexed, high dosage artificial insemination efforts is thus quite significant. The advances achieved by the present inventors in both sexed, unsexed, and low dose artificial insemination represent significant advances which may, for the first time, make commercial applications feasible to equids.

Another problem which has been faced by those in the industry -- again, in spite of the great advances by the Johnson patent and related technology -- is the fact that the problem itself, namely, equine artificial insemination with a high success rate is one of a statistical nature in which a multitude of factors seem to interplay. Thus, the solutions proposed may to some degree involve a combination of factors which, when thoroughly statistically studied, will be shown to be necessary either in isolation or in combination with other factors. Such a determination is further compounded by the fact that the results themselves vary by species and may be difficult to ascertain due to the fact that testing and statistical sampling on a large enough data base is not likely to be worth the effort at the initial stages. For these reasons the invention can also involve a combination of factors which may, individually or in combination, represent the appropriate solutions for a given application. This disclosure is thus to be considered broad enough so that the various combinations and permeations of the techniques disclosed may be achieved. Synergies may exist with other factors. Such factors may range from factors within the sorting, or perhaps, flow cytometer, steps to those in the collection as well as insemination steps. Thus, while there has been a long felt but unsatisfied need for high speed, low dose sexed equine

insemination, and while certain of the implementing arts and elements have long been available, prior to the present invention the advances or perhaps combinations of advances had apparently been overlooked by those skilled in the art. It may even be that the proper combination of known elements simply was not realized. Perhaps to some degree those in the field may have failed to appreciate that the problem involved an interplay of factors as well as peculiar necessities for equine sperm cells involved in this field. Interestingly, as the listing of efforts later in this discussion shows, substantial attempts had been made but they apparently failed to understand the problem inherent in such an area as low dose, sexed insemination of equines and had perhaps assumed that because the natural service event involves perhaps a billion sperm, there may have been physical limitations to the achievement of artificial insemination with numbers which are as many as three orders of magnitude less in number. Thus, it may not be surprising that there was to some extent an actual teaching away from the technical direction in which the present inventors went. Perhaps the results may even be considered unexpected to a degree because they have shown that sexed, low dose equine artificial insemination -- if done right -- can be achieved with success rates comparable to those of unsexed, high dose equine artificial insemination. It might even be surprising to some that the techniques and advances of the present invention in fact combine to achieve the great results shown. While each technique might, in isolation, be viewed by some as unremarkable, in fact, the subtle changes or combination with other techniques appear to afford significant advances in the end result.

Thus, in one regard until the present invention the achievement of non-surgical practical equine artificial insemination low dose, sexed artificial insemination of equines has not been possible with levels of performance necessary or simplified procedures likely to be necessary to achieve commercial implementation. Beyond low dose, sexed insemination on a commercial level, however achieved, the present invention also discloses techniques which permit the achievement of improved performances and thus facilitates the end result desired, namely, low dose, sexed and unsexed non-surgical artificial insemination of equines on a commercial basis.

III. DISCLOSURE OF INVENTION

Accordingly, the invention discloses the achievement of systems for the non-surgical artificial insemination of equine mares. These techniques are applicable for the use of low dosages of equine sperm and are designed so as to be able to be used in the field on a commercial and practical level. Further, the systems are usable in conjunction with -- and have especially valuable applicability to -- artificial insemination with sexed equine sperm. The systems also provide for an improved ability to sort equine sperm cells to determine their sex through flow cytometer separation techniques. Various techniques and substances are represented but as those skilled in the art will readily understand, various combinations and permutations can be used in the manner which may be optimized for performance based on the needs, separation techniques, goals and other parameters involved in a specific equine processing application.

As it relates to the sexed equine application, The objectives of the invention were to 1) compare pregnancy rates in mares inseminated on a single occasion, close to ovulation, with 500, 25, or 5×10^6 progressively motile spermatozoa (pms), 2) achieve reasonable pregnancy rates following insemination with 25×10^6 live-sorted, sexed spermatozoa, and 3) develop techniques for sorting semen.

In one of the initial experiments, sixty-one mares were randomly assigned to 1 of 3 treatments: Group 1 (n=20) were inseminated into the uterine body with 500×10^6 sperm (controls). Group 2 (n=21) and group 3 (n=20) were inseminated in the tip of the uterine horn ipsilateral to the preovulatory follicle with 25, and 5×10^6 sperm. Mares were administered cloprostenol ($250 \mu\text{g}$ i.m.) to induce luteolysis and monitored by ultrasonography every other day until a follicle $\geq 30\text{mm}$ was detected, and then daily until ovulation was detected. GnRH (deslorelin 2.2 mg, Ovuplant®, Fort Dodge) was administered when the dominant follicle was $\geq 35\text{mm}$. Mares were inseminated 34 (n=29) or 40 hours (n=32) after GnRH. Data from 22 mare cycles were excluded because they either ovulated prior to planned insemination

(n=1 1), did not ovulate (n=3), or ovulated > 4 days after GnRH administration (n=8). Semen was collected and immediately diluted with a skim milk extender (EZ-Mixin, OF, Animal Reproduction Systems, Chino, CA) to either 25×10^6 or 5×10^6 motile sperm/ml. Mares receiving 1 ml were inseminated with a flexible plastic artificial insemination pipette (IMV, France), while mares receiving 0.2 ml were inseminated using a disposable implant gun (Veterinary Concepts, Green Valley, WI) containing a 0.5 ml straw. Different insemination pipettes were used to optimize delivery of the two different volumes. The location of pipettes within the uterus was confirmed by transrectal ultrasonography prior to semen deposition.

Pregnancy was determined by ultrasonography at 16 days after ovulation. Pregnancy rates were not different between stallions ($P>0.05$), so results from the two stallions were combined. There was a difference in pregnancy rates for mares bred with 500×10^6 (18/20 = 90%) versus 25×10^6 treatments (12/21 = 57%) ($P<0.05$). There was no difference between mares bred with 25×10^6 versus 5×10^6 treatments (7/20 = 35%) ($P>0.05$). There was no difference in pregnancy rates between mares bred 34 vs. 40 hours after GnRH administration 19/29 (65%) and 18/32 (56%), respectively ($P>0.1$). There was also no difference in pregnancy rates between mares bred with 5×10^6 sperm in a volume of 1 ml, 3/10 (30%) or a volume of 0.2 ml, 4/10 (40%) ($P>0.05$). In summary, pregnancy rates decreased as the number of motile spermatozoa inseminated decreased in this initial effort. However, a day-16 pregnancy rate of 57% was achieved with a single insemination, close to ovulation, with 25×10^6 pms when deposited into the tip of the uterine horn.

In another experiment, seventeen mares were randomly assigned to 1 of 2 treatment groups: Group A (n=1 1) mares were inseminated with approximately 25×10^6 live sorted sperm in a volume of 1 ml. Sperm were sorted into a commercial skim milk semen extender (EZ-Mixin, OF, Animal Reproduction Systems, Chino, CA). One mare failed to ovulate and was excluded from the study. Group B (n=1 0) mares were inseminated with approximately 25×10^6 live sorted sperm in a volume of 1 ml. Sperm were sorted into EZ-Mixin + 4% egg-yolk (EY) Two mares (one from each group) were inseminated with 20×10^6 sperm because of time constraints with the flow cytometers. In both groups, inseminations were performed 34 hr after GnRH administration and sperm were deposited into the tip of the uterine horn,

ipsilateral to the preovulatory follicle using a flexible plastic AI pipette. The location of the pipettes within the uterus was confirmed by transrectal ultrasonography prior to semen deposition. Mares were administered cloprostenol (250 μ g i.m.) to induce luteolysis and monitored by ultrasonography every other day until a follicle ≥ 30 mm was detected, and then daily until ovulation was detected. GnRH (deslorelin 2.2 mg, Ovuplant®, Fort Dodge) was administered when the dominant follicle was ≥ 35 mm. Two stallions were used in this experiment, one of which (Stallion A) was used in Experiment 1. Freshly collected semen was extended 1:1 in HBGM-3 and centrifuged for 10 minutes at 400 x g at 22°C. The supernatant was aspirated and sperm were incubated in 25 μ l Hoechst 33342 at 400 x 10⁶ sperm/ml in HBGM-3 for 1 hr at 35°C and then diluted to 100 x 10⁶ sperm/ml for sorting. Sperm were sorted for sex chromosomes based on a difference in DNA content. Two MoFlo® flow cytometer/cell sorters equipped with an Argon laser emitting 150 mW power at 352 and 364 nm, operating at 50 psi with HBGM-3 as sheath fluid were used for sorting. Aliquots of sorted X and Y populations were reanalyzed for DNA and gave purities of 90 and 84% for X and Y, respectively. Sperm were collected at approximately 900 sperm/sec into 14 ml tubes containing either 4 ml EZ-Mixin (Group A) or 4 ml EZ-Mixin + 4% egg-yolk (Group B). Collected sperm were centrifuged and suspended to 25 x 10⁶ sperm/ml and immediately inseminated. Pregnancy was determined by ultrasonography at 12, 14, 16 and 30 days post-ovulation, and fetuses were sexed 60-70 days post-ovulation without knowledge of the sex of the sorted sperm inseminated. Pregnancy rates were not different between stallions (Stallion A = 3/10, 30%; Stallion B = 5/10, 50%) ($P > 0.1$), so the data sets were combined. Although there was no difference in pregnancy rates between sperm treatments (EZ-Mixin = 3/10, 30% versus 4% EY+ EZ-Mixin = 5/10, 50%) ($P > 0.1$) this may not ultimately prove to be true. At day-60, 5/20 (25%) mares were pregnant; fetuses were sexed and the phenotypic sex ratio was predicted perfectly, five out of five.

This trial has demonstrated for the first time, that pregnancy in the mare can be achieved, and foals of predetermined sex can be obtained, following non-surgical insemination with sexed semen. This is explained in the following discussion. In addition, to the extent they may be helpful, more general sexed insemination aspects and those applicable to equines specifically are discussed in PCT Publication No. _____ by the owner of this

application. Further, the original disclosures of this invention are set forth in United States Patent Applications, serial numbers 60/094,720 and 60/113,143. Each of these prior documents are hereby incorporated by reference for convenience.

5 Thus, an object of the invention is thus to achieve artificial insemination of equines in a field environment with no need to resort to surgical procedures. Further, a goal is to provide the ability to use lower dosages in a manner which works under realistic commercial circumstances and which yields pregnancy success probabilities which are comparable to traditional equine dosage success rates. An object is also to achieve better sorting for equine sperm cells. A parallel goal is to provide substances and techniques which are especially
10 suited for equine sperm cells when being separated into X- and Y-chromosome bearing components. Thus a goal is to achieve a sorted result which is consistent with unsorted, high dosage expectations.

A goal is also to present an overall system for equine artificial insemination which can achieve its objects in a commercially practical manner. Sorting in a manner which affords
15 both high speed and low stress equine sorting, and which is especially adapted for equine sperm cell sorting in a low dose context is an important goal as well.

Naturally further objects of the invention are disclosed throughout other areas of the specification and claims.

IV. BRIEF DESCRIPTION OF DRAWINGS

20 Figure 1 is a diagram of a sorter system according to a flow cytometer separation technique for the present invention.

Figure 2 is a diagram of the entrained cells in the nozzle just prior to the free fall area of a typical flow cytometer.

V. BEST MODE FOR CARRYING OUT THE INVENTION

As will be seen, the basic concepts of the present invention can be combined and embodied in a variety of ways. The invention involves commercially practical low dose, sexed equine artificial insemination and the results. For flow cytometry separation techniques, the invention also involves both improved flow cytometer systems as well as systems for the creation of sex-specific equine sperm samples which may be used in equine artificial insemination and the equids produced by such techniques. It discloses overall processes through which high success rates are possible even in commercial equine environments. Furthermore, the techniques are disclosed in a general fashion so that they may be applied to specific systems and applications once the general principals are understood. While device enhancements are disclosed it should be understood that these enhancements not only accomplish certain methods but also can be varied and combined in a number of ways. Importantly, as to all of the foregoing, each of these facets should be understood to be encompassed by this disclosure.

When considering the sex selection aspect of the invention, the basic goal is that of separating the X-bearing sperm from the Y-bearing equine sperm in a way that then can be used to artificially inseminate the mare with high success rates. Preferably this insemination would not require surgery. The separation phase is preferably done in a manner which isolates the two types of equine sperm so that each can be separately packaged and dealt with. At present the isolation is preferably done through the use of flow cytometry. Flow cytometry in general is a technique which is well understood. For instance, the basic aspects of it are shown and discussed in a variety of patents to Cytomation, Inc. such as the U.S. Patents and other publications listed earlier. In addition, some details as applicable to equines and other species are disclosed in PCT Publication No. _____. Each of these publications and the references cited therein, are incorporated by reference so those skilled in the art can easily understand the basic principles involved.

In general, it should be understood that during meiosis in the testis, the sex chromosomes segregate into individual spermatids and haploid spermatozoa carry either the

X or Y chromosome. There is a 50:50 ratio of X- to Y- bearing spermatozoa in the semen, and fertilization of an X-bearing, haploid oocyte by either an X-or Y-bearing sperm determines the sex of the embryo. A 50:50 ratio exists because X- and Y-bearing spermatozoa are made in equal numbers and are phenotypically identical. The desire to alter this 50:50 ratio and predetermine the sex of mammalian offspring has been of great interest to the public for many years. There are numerous benefits to sex preselection in equine animals. Sex preselection has also been used to produce females when heritable X-linked diseases are an issue. Unlike humans and most farm animals, the advantage of sex preselection in the horse can also be purely one of preference to the breeder/owner and there has been considerable interest expressed by members of certain breed registries.

For years, numerous attempts to separate X- from Y- chromosome bearing spermatozoa have been made based on physical and chemical properties of sperm. Johnson (referenced earlier) tested these methods and found that the only method proven effective was based on a difference in DNA content of the spermatozoa. No other method based on a physical difference within the spermatozoa or on surface properties has been proven effective in separating X- from Y-bearing spermatozoa. Within equines, DNA content of mammalian X- and Y-chromosome bearing sperm differs by 4.1%. This difference in DNA content can be used to separate X- and Y-chromosome bearing spermatozoa after staining sperm with a fluorescing, DNA binding dye followed by flow cytometry.

Modern flow cytometry/cell sorting technology was first developed by Fulwyler in 1965. Flow cytometry has mainly been used in medical research and diagnoses with respect to blood and tumor cells, but can also be used to evaluate many types of cell suspensions including sperm cells. Essentially, flow cytometry as applied here involves sorting equine sperm cells, which are provided to the flow cytometer instrument through some type of cell source. A conceptual instrument is shown in Figure 1. The flow cytometer instrument includes a sample input, here an equine sperm cell source (1) which acts to establish or supply equine sperm cells or some other type of item to be analyzed by the flow cytometer. The cells are deposited within a nozzle (2) in a manner such that the cells are surrounded by a sheath fluid (3). The sheath fluid (3) is usually supplied by some sheath fluid source (4) so that as

the equine sperm cell source (1) supplies its cells, the sheath fluid (3) is concurrently fed through the nozzle (2). In this manner it can be easily understood how the sheath fluid (3) forms a sheath fluid environment for the equine sperm cells. Since the various fluids are provided to the flow cytometer at some pressure, they flow out of nozzle (2) and exit at the nozzle orifice (5). By providing some type of oscillator (6) which may be very precisely controlled through an oscillator control, pressure waves may be established within the nozzle (2) and transmitted to the fluids exiting the nozzle (2) at nozzle orifice (5). Since the oscillator (6) thus acts upon the sheath fluid (3), the stream (7) exiting the nozzle orifice (5) eventually and regularly forms drops (8). Because the cells are surrounded by a sheath fluid environment, the drops (8) may contain within them individually isolated cells or other items.

Since the drops (8) generally contain isolated equine sperm cells, the flow cytometer can distinguish and separate droplets based upon whether or not the appropriate cell or cells is/are contained within the drop. This is accomplished through a cell sensing system (9). The cell sensing system involves at least some type of detector (10) (which may include two detectors at 90 degrees with respect to each other) which responds to the cells contained within each drop (8) as discussed at length in the seminal work (no pun intended) by Larry Johnson, namely, U.S. Patent No. 5135759. As the Johnson patent explains for sperm cells, the cell sensing system (9) may cause an action depending upon the relative presence or relative absence of a particular dye which may be excited by some stimulant such as the laser exciter (11). While each type of sperm cell is stained by the dye, the differing length of the X-chromosome and the Y-chromosome causes different levels of staining. Thus, by sensing the degree of dye present in the sperm cells it is possible to discriminate between X-bearing sperm and Y-bearing sperm by their differing emission levels.

In order to achieve the ultimate separation and isolation of the appropriate cells in a flow cytometer separation technique, the signals received by detector (10) are fed to some type of sorter discrimination system (12) which very rapidly makes the decision and can differentially charge each drop (8) based upon whether it has decided that the desired equine sperm cell does or does not exist within that drop (8). In this manner the sorter discrimination system (12) acts to permit the electrostatic deflection plates (13) to deflect

drops (8) based on whether or not they contain the appropriate cell or other item. As a result, the flow cytometer acts to sort the cells by causing them to land in one or more collectors (14). Thus by sensing some property of the cells or other items the flow cytometer can discriminate between equine sperm cells based on a particular characteristic and place them in the appropriate collector (14). In the system presently used to sort equine sperm, the X-bearing sperm droplets are charged positively and thus deflect in one direction, the Y-bearing sperm droplets are charged negatively and thus deflect the other way, and the wasted stream (that is unsortable cells) is uncharged and thus is collected in an undeflected stream into a suction tube or the like.

Referring to Figure 2, the process can be even further understood. As shown in that figure, the nozzle (2) emits a stream (7) which because of the oscillator (6) (not shown in Figure 2) forms drops (8) (not shown in figure 2). Since the equine sperm cell source (1) (not shown in Figure 2) may supply equine sperm cells (15) which have been stained according to the Johnson technique, the light stimulation by laser exciter (11) and differentially determined state as sensed by detector (10) can be used to create the existence or nonexistence of a charge on each drop (8) as it separates from stream (7). This is all controlled by the flow cytometer. This control results in positively charged, negatively charged, and uncharged drops (8) based upon their content. As shown in Figure 1, certain drops are shown as deflected drops (16). These deflected drops (16) are those containing sperm cells (15) of one or the other sex. They are then deposited in the appropriate collector (14) for later use.

As shown in Figure 2, the equine sperm cells (15) may be injected into the sheath fluid (3) by a needle or injection tube (18). As those skilled in the flow cytometry arts understand, this injection tube (18) can be shaped so as to achieve hydrodynamic focusing and orientation so that both the tail and/or the flatter side of the equine sperm cells (15) are properly oriented. It may even produce a ribbon shaped sample core stream as shown. This can be important since the ninety degree fluorescence intensity can be viewed as proportional to sperm head orientation and the zero degree fluorescence intensity can be viewed as proportional to the sperm DNA content.

High resolution flow cytometric DNA analysis of equine sperm is difficult compared to other cells because the highly compacted chromatin in the morphologically flat, paddle shaped sperm head, causes a high index of refraction. The difference in refractive index between the sperm head and the surrounding medium, in combination with the flat shape of the sperm head, results in more emission of fluorescence through the plane of the cell (from the edge of the sperm head) than at a 90° angle to the plane. Proper orientation of the head is critical for high resolution sorting. Others have investigated methods of controlling the orientation of sperm heads flowing in a stream and found that a beveled injected tube (18) was effective in subjecting cells to planar hydrophobic forces as the cells flowed out the end of the needle. The sample stream can be pushed into a ribbon shaped stream which can orient the flat head of the sperm in the same manner. Other, more complex, physical principles are often applied to orient sperm as well.

As mentioned, spermatozoa that undergo the sorting process are stained with the fluorochrome stain Hoechst 33342, chemically known as bisbenzimidazole. The dye is added to the sample at approximately 9 μ M in a 1 ml volume containing 400 x 10⁶ sperm, and may be incubated at 32-35°C. Hoechst 33342 is non-toxic to sperm, and does not significantly alter sperm motility. Hoechst 33342 binds to the A-T regions of the DNA helix. Since the sorting process is more productive if dead sperm are not collected, a molecule which quenches the Hoechst 33342 fluorescence and marks dead sperm can be added to the stained sample. Food coloring is one example of a nontoxic molecule that has been used. The process is then that fluorescently stained spermatozoa are introduced under pressure (e.g. 50 psi) in liquid suspension to the flow cytometer. The sperm enter the sample insertion tube and are oriented in one way or another, perhaps as they exit the beveled end of the tube into sheath fluid. The stream containing the sperm intersects an Argon-ion laser beam, in ultraviolet (351 and 364 nm) wavelengths at up to 200 mW of power. Approximately 7% of the drops contain a sperm cell when the sample is at 100 x 10⁶ sperm/ml. The fluorescently stained nuclei are excited by the laser beam, giving off fluorescent signals proportional to the amount of DNA bound to the dye.

A modified commercial flow sorting system was used to analyze and sort spermatozoa based on DNA differences. As mentioned, the fluorescent signal from the sperm edge (90° angle from laser emission) is brighter than that emitted from the flat side (0° forward detector). The edge emission is used to characterize the orientation of the sperm heads as they pass the laser beam. The emitting light is collected by the 90° detector, and properly oriented sperm are recognized. Sperm that are not oriented properly give off less light and are electronically gated out of the analyses often, because of the lack of proper orientation, the majority of sperm are not separated into X or Y enriched populations. Also, many sperm cells that are properly oriented are not sorted because of the distributions of fluorescence overlap. An optical detector and a photomultiplier tube (PMT) can be used for the 0° detection of fluorescence, which is proportional to the DNA content of properly oriented sperm. The PMT's collect the fluorescence emitted by the sperm and convert the optical signal into a proportional electronic signal, which is amplified by the PMT for further signal processing and graphic display. Electronic gating enables selection of signals from the 0° forward detector only for properly oriented sperm; this results in the ability to differentiate between the DNA of X and Y sperm. By setting the electronic sort windows around the resolved populations, X- and Y-bearing sperm are separated into two tubes. Two optical detectors collect the light given off, and circuits are activated which add a charging pulse (+ or -) to the drops containing the respective X- or Y-bearing sperm. As the individual sperm droplets fall, they pass an electrostatic field that pulls the charged droplets containing sperm into separate tubes which contain "catch fluid" extender as a temporary holding medium until the sperm are further processed.

Spermatozoa samples which have been separated into X- and Y-chromosome-enriched samples can be reevaluated for their purity using flow cytometric analysis for the DNA content of the individual sperm. In the past, the only valid method for verification of separation available was to determine the sex ratio of the offspring which is untimely and expensive. Re-analyses using flow cytometry reduces not only time and cost, but increases the precision.

With advances it is anticipated that the percent of sperm that are oriented properly as the droplets pass the laser can increase, resulting in increased sorting rates from 100 live sperm/s of each sex to rates between 1000 and 1500 live sperm/s of each sex at ~ 90%. This increase in the sperm sorting rate would be helpful to make it more practical to utilize sexed semen in an equine artificial insemination program.

5

Others have reported using the amount of DNA in sperm as a marker for sex preselection and subsequent birth in rabbits of the predicted sex following surgical insemination. Spermatozoa were separated into X- and Y-chromosome bearing populations with a flow cytometer/cell sorter. Sorted sperm were surgically inseminated into the uterus. From does inseminated with fractions enriched for Y-bearing sperm, 81% of offspring born were male; fractions enriched for X-bearing sperm resulted in 94% females. Thus the sex ratio was significantly altered from 50:50. This phenotypic (and genotypic) ratio was accurately predicted based on the reanalysis for DNA from the sorted populations, which showed purities of 81% for Y-bearing sperm and 86% for X-bearing sperm. This was followed by publication of experiments in swine, and cattle. *In vitro* fertilization has been used to obtain pregnancies in cattle using sexed semen. For example, one researcher transferred twin embryos into each of 9 heifers. Four heifers became pregnant and 6 calves were born, all of the predicted sex. These results show that viable sperm can be separated into X- and Y-chromosome bearing populations and retain their capacity for fertilization and producing normal progeny. Pregnancies following *in vitro* fertilization (IVF) with X- and Y-bearing sperm have also been obtained in swine and rabbits. Recently, bovine embryos have been produced by IVF with sperm sorted by high speed flow cytometry, but embryo transfers were not done.

10

15

20

Using sorted-sexed equine sperm in a laboratory environment, one researcher performed two experiments to determine pregnancy rates with intracytoplasmic sperm injection (ICSI) into equine oocytes or oviductal insemination. Thirteen injected oocytes developed to the 2- to 3-cell stage, 8 to the 4- to 6 cell stage and 2 oocytes developed to the 7- to 8-cell stage. One embryo was transferred at the 7- to 8- cell stage, but no pregnancy resulted. In the oviductal insemination experiment, two mares were inseminated by

25

cannulation of the fimbriated end of the oviduct with 50 μ l containing 1.5×10^5 sorted X-bearing sperm. One pregnancy was detected and the mare produced a female foal.

One of the aspects of flow cytometry which is particularly important to its application for equine sperm sorting is the high speed operation of a flow cytometer. Advances have been particularly made by the flow cytometers available through Cytomation, Inc. under the MoFlo® trademark. These flow cytometers have increased sorting speeds extraordinarily and have thus made flow cytometry a technique which is likely to make feasible the commercial application of equine sperm sorting (among other commercial applications). They act to achieve high speed sorting, that is at a speed which is notably higher than those otherwise utilized. Specifically, Cytomation's MoFlo® flow cytometers act with oscillator frequencies of greater than about five kilohertz and more specifically can be operated in the 10 to 30 or even the 50 kilohertz ranges. Thus droplets are formed at very high frequencies and the cells contained within the sheath fluid environment can be emitted very rapidly from the nozzle (2). As a result, each of the components such as the nozzle (2) oscillator (6), and the like which make up and are part of a flow cytometer system can be configured or selected to result in a high speed cell sorter. In the application of a high speed cell sorter to the sorting of sperm cells, sorting at rates of greater than about 900 sorts per second is achieved. Importantly, it should be understood that the term "high speed" is a relative term such that as other advances in flow cytometry and specific applications are achieved, the aspect which is considered "high" may be varied or may remain absolute. In either definition, the general principle is that the sorting may occur at rates at which the parameters and physical characteristics of the flow cytometer are significant to the cells themselves when sorting particular cells such as equine sperm cells.

One aspect of high speed sorting which appears to come into play when sorting equine sperm cells through a flow cytometer separation technique is that of the pressures and other stresses to which the equine sperm cells are subjected within the flow cytometer. For instance, when operating at high speeds (and an alternative definition of "high speed"), flow cytometers can be operated at a pressure of 50 pounds per square inch and even higher. These pressures may be considered high because they may result in effects upon the equine

sperm cells being sorted. The key as disclosed in the present invention for this facet is the fact that the stress thresholds of the particular cells are the determining factor. Additionally as further knowledge is gained it may be shown that the stress thresholds are a function of combined effects such as the particular species or the particular prior or subsequent handling of the equine sperm cells. The key in this regard is that the stress imposed upon the equine sperm cells can, in fact, alter their viability and their ability to achieve the desired result. This may be unusually true for equine species. In the pressure case, it may be that merely subjecting the sperm cells to a higher pressure as a result of the operation of the flow cytometer at that pressure may result in decreased performance of the equine sperm cells. The present invention in one regard acts to minimize these stresses and thus results in greater efficacies as well as lower dosages as discussed later.

In considering the stress aspect of the equine cells, the present invention acts in a fashion which minimizes the stresses. These stresses can be minimized at any point in the over-all cycle or process of collecting, sorting or even inseminating the animal. Importantly, the stress imposed by the handling of the cells within the flow cytometer appears significant for equine species. In one embodiment of the invention, the sheath fluid is specifically selected so that it can serve in a coordinated fashion with both (or either) the pre-sort cell fluid environment or the post-sort cell fluid environment. Thus the present invention acts to minimize the changes through the type of operation or the selection of substances which may act as a means for minimizing the changes which the equine sperm cells experience. For the sheath fluid, a substance is selected according to one embodiment of the invention so that it may be chemically coordinated to present minimal changes. Thus, by selecting the appropriate sheath fluid not only in context of flow cytometry parameters, but rather also in context of the equine sperm cell and equine artificial insemination parameters themselves, the changes experienced by the cells and the over all result of the sorting can be enhanced. Interestingly, for equine sperm cells, it has been discovered that a hepes buffered medium such as a hepes bovine gamete medium — particularly HBGM3 as previously created by J. J. Parrish for a bovine application — works well. This medium is discussed in the article "Capacitation of Bovine Sperm by Heparin", 38 Biology of Reproduction 1171 (1988) (hereby incorporated by reference). Not only is this surprising because it is not the same type

of use as for bovine sperm, but the actual buffer, was originally developed for a bovine application. Thus in the equine application the sheath fluid is selected which contains the hepes buffer.

5 A separate aspect of the flow cytometer processing which may also be important is the fact of properly treating the cells both chemically and physically after they are sorted. As shown in figure 2, the cells within drops (8) land in collector (14). The collector fluid (17) can also serve to minimize stresses upon the cells. In one regard, since it may be important to provide a nutrient to the cells both before and after sorting, the collector fluid (17) may be selected so as to provide an easier reception as well. For equine sperm cells, the cells may be
10 collected in a commercial skim milk extender such as that from EZ-Mixin, Animal Reproduction Systems, Chino, CA. Even though intended for a different purpose, this extender can be used as the collection fluid for equine sperm cells. Further, this collection fluid may include 4% egg yolk as well.

15 Another aspect which may interplay in the various factors of the present invention is that of utilizing low dose amounts of sperm for artificial insemination or the like. Additional background on the aspect of sexed, artificial insemination may be found in "Prospects for Sorting Mammalian Sperm" by Rupert P. Amman and George E. Seidel, Jr., Colorado Associated University Press (1982) and in the previously referenced PCT publication, each hereby incorporated by reference. As mentioned earlier, natural insemination in equines
20 involves numbers of sperm on the order of a billion of sperm. Routine artificial insemination is presently conducted with two hundred-fifty million or more sperm for equine species. By the term "low dose" it is meant that the dosage of sperm utilized in the insemination event are less than one-quarter or preferably even less than about 10% or 5% of the typical number of sperm provided in a typical equine artificial insemination event. Thus, the term "low dose" is
25 to be viewed in the context of the typical artificial insemination dosage or also as an absolute number. The absolute numbers may be species dependent, of course. For equine species, merely less than about twenty-five, ten, five, or even one million sperm may be considered a low dose process.

Yet another aspect which may be important is the fact that the sperm sexed through the present invention techniques (or otherwise) is utilized in an equine artificial insemination system. Thus when, for a flow cytometer technique, the collector (14) is used to provide sperm for artificial insemination the techniques of the present invention may be particularly relevant. Further, it is possible that the combination of both equine artificial insemination use and the use in a low dose environment may together create synergies which makes the various techniques of the present invention particularly appropriate. Naturally, the sexed sperm can be utilized not just in an artificial insemination mode, but in other techniques such as in vitro fertilization and the like.

The process of collecting, sorting, and eventually inseminating an animal through the use of a flow cytometry sorting, or other separation technique, involves a variety of steps. In the context of equine insemination, first the semen is collected from the stallion. Semen may be collected from stallions of known high fertility immediately prior to planned insemination. This may occur with a Colorado model artificial vagina (Animal Reproduction Systems, Chino, CA) equipped with an in-line gel filter. Ejaculates can then be evaluated for gel free-volume, motility and spermatozoal concentration. Semen can be then extended with a commercial skim-milk glucose extender (EZ-Mixin, OF, Animal Reproduction Systems, Chino, CA) to either 25×10^6 pms/ml ($n=51$) or 5×10^6 pms/ml ($n=10$). Semen may be kept at room temperature until inseminations were performed, shortly after collection. Staining may be accomplished according to a multi-stained or single-stained protocol, the latter, the subject of the Johnson Patent and related technology.

After adding the stain, dilution or extending to the desired sort concentration may be accomplished. Sorting according to the various techniques discussed earlier may then be accomplished from which sperm cells may be recovered in the collection phase.

An optimal number of motile spermatozoa per insemination dose to maximize fertility with prior techniques has perhaps been well established in species such as swine, sheep, and cattle. With the present invention, the minimum number of motile spermatozoa seem to be much less than the 250 to 500×10^6 progressively motile spermatozoa (pms) usually

recommended. Under ideal conditions, mares had even been inseminated with as few as 100×10^6 pms without reducing fertility. (Researches found no difference between mares inseminated over three cycles with 100 or 500×10^6 pms and achieved pregnancy rates of 63.9 and 75%, respectively.) The present invention shows even lower numbers now to be possible -- and that the lower numbers can be achieved in a field environment.

Although the difference was not significant, in one experiment pregnancy rates for the 100 and 500×10^6 treatments for cycles 1, 2, and 3 were 25 vs. 39, 33 vs. 45, and 28 vs. 25% respectively. Notably, others have reported an increase in foaling rate when the number of motile spermatozoa per insemination was increased from 40 to 80×10^6 , but no further improvement was observed when the number of spermatozoa was increased to 160×10^6 . In an experiment using two groups of 14 subfertile mares, one researcher found no difference between treatments utilizing 100 or 500×10^6 motile spermatozoa per insemination (35.7 vs. 42.9%, respectively). Later, the same researcher reported pregnancy rates after breeding over three cycles from mares inseminated with 50, 100, and 500×10^6 pms of 41.7, 65.6, and 81.3%, respectively, from data averaged from several experiments. Yet another in the art inseminated mares (over three cycles) with 50 and 500×10^6 pms and found a significant difference between pregnancy rates of 37.5 and 75%, respectively. In a more recent study, one of the inventors superovulated mares with equine pituitary extract (EPE) and inseminated mares one time with 50×10^6 pms. Pregnancy rates were not different between the mares treated with EPE and the saline controls (65 and 55%, respectively).

In existing routine equine artificial insemination, though there appears to be little difference in fertility between 100 and 500×10^6 spermatozoa per insemination dose, 500×10^6 pms is generally recommended to provide maximum fertility. However, when proper artificial insemination techniques were utilized, 100×10^6 pms from a highly fertile stallion was also believed to be the minimum adequate with the prior techniques. In the more accepted circumstance, when performing routine artificial insemination (AI) with fertile stallions and mares, 500×10^6 progressively motile sperm/dose inseminated every other day while mares are in estrus has been reported to result in maximum fertility. As alluded to

earlier, the problem with this is simple, insemination of mares with a low number of spermatozoa may be necessary when semen is limited or when using sorted sexed semen.

As mentioned earlier, currently, it is possible to obtain approximately 1000 equine live sperm/second ($3.6 \times 10^6/\text{hr}$) of each sex chromosomal composition when sorting spermatozoa for sex chromosomes by flow cytometry at 90% accuracy. Thus it would be impractical to obtain 500×10^6 sperm that were sorted for sex chromosomes for an insemination at 3.6×10^6 sperm/hour. The goal therefore, was to achieve lesser numbers of spermatozoa per insemination dose while obtaining reasonable fertility. The objective was to achieve pregnancy rates in mares inseminated on a single occasion, close to ovulation, with as low as 25 or 5×10^6 or lower progressively motile spermatozoa (pms).

The normal stallion ejaculate contains an average volume of 50 ml. The stallion deposits this high volume of semen directly into the uterus of the female. In another species (boar) which ejaculates several hundred ml of semen, only 0.5-0.1 ml enters each fallopian tube at the beginning of estrus to permit fertilization. This large seminal volume fills the region of the utero-tubal junction until a reservoir of spermatozoa is established in the isthmus. Therefore a specific volume of semen is established in the isthmus and the remaining content is rapidly eliminated.

For artificial insemination, the number of spermatozoa in an insemination dose can be critical. Seminal extenders are often used to dilute raw semen to provide larger and more easily managed insemination volumes. In the prior art, volume ranging between 10 and 25 ml of semen is generally recommended although, now, perhaps depending on the concentration of spermatozoa, small insemination volumes can prove to be as effective as larger volumes. Although the concentration was not specified, insemination volumes ranging from 0.6-26.8 ml of semen did not adversely affect fertility when the effects of inseminating 10 or 50 ml volumes of extended semen on embryo recovery rates were studied in mares. But based on this experiment, there was a reduction in embryo recovery rates from mares inseminated with a 50 ml volume of extended semen compared to a 10 ml volume when both contained an equal number of pms. The reduced fertility may have been due to the increased volume

inseminated or could have been due to the decreased spermatozoa concentration since it was $1/5^{\text{th}}$ that of the 10 ml volume (5 vs. 25×10^6).

Further experiments were conducted to determine: 1) embryo recovery rate when mares were inseminated with 100×10^6 pms extended in 10, 100, or 200 ml of dried skim milk extender and 2) embryo recovery rate when mares were inseminated with 250×10^6 pms extended either 10 or 100 ml of the same extender as experiment 1. Results from experiment 1 showed a difference in embryos recovered only between mares inseminated with 10 (40%) and 200 ml (0%). In one of those experiments there was a significant difference between mares inseminated with 10 ml (70.6%) compared to 100 ml (13%). Thus, insemination volumes of 100 or 200 ml were associated with lower embryo recovery rates than a 10 ml volume, probably due to the lower sperm concentration or retrograde loss of sperm into the vagina.

They conducted a study to test whether volume alone affects fertility when sufficient concentrations and numbers of spermatozoa are present. They concluded that there was no difference between mares inseminated with either 30 or 120 ml of cooled semen at a concentration of 50×10^6 pms/ml. This approach, however, is not followed by the present invention to some degree.

Timing and frequency of insemination can play a very important role in most breeding operations, especially when frozen or shipped-cooled semen is involved. The number and timing of inseminations can affect fertility. The average mare ovulates every 21 days during the physiological breeding season, and the average duration of estrus during this time is 5-7 days. During estrus, mares will passively urinate, lift their tail, and present their hindquarters to the stallion. Under natural conditions when a stallion was introduced to a herd of 20 mares, the number of breeding per hour of observation was 2.4 ± 0.2 . Stallions often breed the same mare multiple times per day (under natural conditions). In one study, 20 mares were synchronized and placed in a pasture with a stallion and observed for 9 days. The stallion mated 9.12 times per day and settled 17 of 18 mares.

Researchers have also compiled data over multiple breeding seasons comparing the effect of the number of inseminations on pregnancy rates. More mares became pregnant when inseminated five times (68%) more than mares that were inseminated three times (35.9%) during cycle 1. No other differences were noted in regard to number of inseminations on pregnancy rates during cycle 1. There was no difference in pregnancy rates during cycles 2 and 3 when mares were inseminated one to seven times. More mares became pregnant when inseminated 5 times (60%) than mares inseminated 1 (23.5%), 2 (35%), or 3 times (35.5%) over 3 cycles. When considering all 3 cycles, mares that became pregnant were inseminated an average of 3.3 times, which was more than the average of 2.8 times for mares not becoming pregnant.

It has also been determined that multiple inseminations per cycle were not detrimental to fertility. In one study, data were collected from 257 mares over a 10-year period to establish the relationship between the number of inseminations per cycle, duration of estrus and pregnancy rates. Mares were inseminated with 100×10^6 spermatozoa. First cycle pregnancy rates of 22.0, 23.0, 38.6, 52.5, 58.3, and 52.2% were achieved when mares were inseminated 1, 2, 3, 4, 5, or 6 or more times per cycle, respectively. Fewer mares became pregnant after three cycles when inseminated 1-4 times per cycle than mares inseminated ≥ 12 times per cycle. Another study inseminated 62 mares over three cycles every 48 hours during estrus with 200×10^6 pms for a maximum of three inseminations. Inseminations began when a follicle ≥ 30 mm was detected and continued until ovulation. Fertility per cycle was 45% and was not different if two or more inseminations were done per cycle as compared to one insemination per cycle. They also determined that the highest pregnancy rates were achieved with inseminations performed between 48 and 72 (8/23) or 72 and 96 hours (8/23) before ovulation and that the last insemination was not the fertilizing one at least 51% of the time. Overall, when performing routine AI with fertile stallions and mares, 500×10^6 pms/dose inseminated every other day while mares are in estrus results in maximum fertility. With the present invention, much lower numbers are now possible.

Induction of ovulation at a specific time in the mare may be advantageous for the following reasons: to ensure that ovulation will occur within 36-48 hours of mating

eliminating the need for rebreeding, (b) with use of cooled, frozen or sexed semen when timing is critical in order to maximize fertility, (c) to ensure that only a single insemination close to ovulation is needed when utilizing subfertile stallions or mares, (d) to minimize mare or stallion transport, and (e) to stagger ovulations when multiple mares are presented in estrus at the same time.

Human chorionic gonadotropin (hCG) is produced by the cytotrophoblast of the chorionic villi of the human placenta. It is a glycoprotein hormone composed of two subunits (α and β) which are linked together non-covalently. It has a half-life of 8-12 hours in blood.

Use of hCG for induction of ovulation during the estrous cycle of the mare was first reported in 1937 by Mirskaja and Petropavlovski. They found that ovulation occurred within 24 to 48 hours after injection of crude extract of human pregnancy urine (Prolan®) injected on the first day of estrous. Further studies have shown that when hCG (1500-3300 IU) is injected in a mare during early estrus, it mimics lutenizing hormone (LH) activity and induces ovulation, generally within 24-48 hours. The use of hCG at a dose of 2000- 3000 IU has not decreased fertility. However, some researchers did find that higher doses (4500-6000 IU) resulted in reproductive disorders and a decreased pregnancy rate. Although the use of hCG can be very effective in inducing ovulation, several researchers have shown that administration of hCG over several consecutive estrous cycles can result in antibody formation, with mean duration of estrus and ovulation either the same as the control mares or 2 days longer than controls.

Native gonadotropin releasing hormone (GnRH) is a decapeptide synthesized in the hypothalamus and stored in secretory granules of the median eminence. Upon release, GnRH enters the portal system and is transported to the anterior pituitary and binds to receptors on gonadotrope cells where it stimulates synthesis and secretion of lutenizing hormone (LH) and follicle stimulating hormone (FSH). Research has also been conducted on the use of native GnRH and GnRH analogues in which 1 or 2 amino acids have been modified on inducing ovulation during estrus in the mare.

Pulsatile or continuous administration of native GnRH causes predictable ovulation. In one study 11 cycling mares were infused with either saline or 20 μ g GnRH in a pulsatile pattern (one 5-sec. pulse/hr, 2h or 4h) starting on day 16 of the estrous cycle. The number of days from start of treatment to ovulation was less in mares infused with 20 μ g GnRH/hr compared to saline control mares or 20 μ g GnRH per 4hr. It was concluded that pulsatile infusion of GnRH is effective in advancing ovulation, but the frequency of the pulse is a critical variable. Native GnRH has also been used to induce follicular development and ovulation in seasonally anestrus mares. A short term implant which releases 1.5 or 2.2 mg of the GnRH analogue deslorelin causes ovulation within 36-48 hours when administered to mares in estrus with a follicle >30 mm in diameter.

One of the inventors has compared the effect of various doses of a GnRH analog (deslorelin acetate) implant, on induction of ovulation in cyclic mares and found that ovulation was induced in most mares within 48 hr after injection and there is no advantage of doses higher than 2.2 mg/mare. Others have compared the use of hCG, buslerelin (a GnRH analog) and luprostitol (a PGF₂ α analog) for induction of ovulation in cycling mares. Both buslerelin and hCG shortened the interval from treatment to ovulation, whereas luprostitol failed to hasten ovulation.

Equine pituitary extract (EPE) is derived from equine anterior pituitary glands. Preparation of EPE for experimentation as a crude gonadotropin has been described by Braselton and McShan (1970), and more recently by Guillou and Combarnous (1983), each hereby incorporated by reference. EPE has been used in the mare primarily to induce growth of multiple follicles in cyclic or anestrus mares and for superovulation in the ewe.

Use of equine pituitary extract as an ovulatory agent in the mare has been known. In studies, some have separated equine luteinizing hormone (eLH) and follicle stimulating hormone (eFSH) by hydrophobic interaction chromatography (HIC) and conducted experiments. In one experiment, LH activity in crude equine gonadotropin (CEG) was compared to LH activity in the HIC fraction on its ability to induce ovulation. Of 25 control mares, 7 ovulated within 48 h compared with 24/25 mares treated with CEG and 19/26 mares

5 treated with LH. Another experiment was designed to test the ability of the eFSH-enriched fraction of pituitary extract to induce the growth of multiple follicles compared to CEG. The number of follicles that reached 30 mm was the same in CEG vs. FSH treated groups and both groups were different when compared to the control group. Ovulation rates were not different between the two treatment groups but were different from the control group.

10 Historically, the most commonly used method of inducing ovulation is a single injection of hCG. This still remains the most common method. However, since there is no difference in pregnancy rates or timing of ovulation when administering either GnRH or hCG to cycling mares, either treatment is an acceptable method for inducing ovulation. EPE however, is not commercially available to practitioners and therefore is not a practical technique for inducing ovulation.

15 Under natural mating conditions, the equine ejaculate is deposited directly into uterus of the mare. Spallanzani was first to report artificial insemination (AI) in dogs, and then horses in the late 1700s. The use of AI has been documented in cattle, sheep, swine, and horses. Horses, cattle, and hogs are artificially inseminated within the uterine body; sheep, goats and dogs in the cervix; and cats in the anterior vagina. As to equines, others have described routine seminal collection and handling procedures in the horse. For routine AI procedures, semen is deposited within the uterine body using a sterile insemination pipette and syringe. However, there are several reasons for the use of alternative sites and techniques for artificial insemination: a) insemination of frozen thawed semen of low quality or limited quantity, b) insemination of semen from a subfertile sire or c) insemination with sexed semen, which is of limited quantity. Some alternative AI techniques include: intra-uterine insemination (via laparoscopy or nonsurgical techniques) in those species where cervical or vaginal inseminations are routinely performed, oviductal insemination (via laparoscopy or flank laparotomy), or deep intra-uterine nonsurgical insemination.

25 Laparoscopic intra-utene insemination has evolved as the least invasive technique for depositing semen directly into the uterus of sheep and goats since the early 1970's when suitable equipment was developed. Laparoscopic insemination is routinely performed in the

ewe and goat with high fertility compared to traditional AI. Laparoscopic intra-uterine insemination has also been successfully reported in the ferret, domestic cat, tiger, cheetah and leopard, and most recently the possum and wallaby. Some advantages of laparoscopic insemination include: genetic improvement utilizing frozen semen, increased number of
5 inseminations per collection using lower sperm numbers, and higher fertility. The main disadvantage of laparoscopic insemination is the higher cost of the equipment and procedure (skilled labor, drugs, semen processing). This procedure is also relatively invasive to the patient.

Nonsurgical intra-uterine insemination with ewes is used in an attempt to increase
10 fertility rates in species that are routinely inseminated in the cervix. One researcher obtained a 75% lambing rate following intra-uterine insemination, compared to 17% after deep cervical insemination, and 30% after double caudocervical insemination. In another study, a researcher deposited frozen-thawed ram sperm into three regions of the genital tract of ewes. In group 1, a single intra-uterine insemination was performed, while in group 2, ewes were
15 inseminated once deep in the cervix, and in group 3, ewes were inseminated twice, 12 hours apart, in the caudocervical region; conception rates were 89, 45, and 57% respectively. Others reported similar results with intra-uterine insemination. Nonsurgical endoscopic insemination has also been performed in bitches resulting in high pregnancy rates.

With oviductal insemination (OI) a small volume of semen (usually 0.05-0.5 ml) is
20 surgically inseminated into the oviductal lumen. One study inseminated nine gilts using laparoscopic insemination. Two of the nine (22%) gilts became pregnant from a single insemination. A more recent study in the ewe, determined the effects of number of spermatozoa, timing and site of insemination on fertility. In experiment 1, ewes were
25 inseminated with 10^4 , 10^5 , 10^6 or 10^7 spermatozoa. Ova recovered 48 hours later were classified as fertilized if they had cleaved. Results showed that more ewes were fertile after oviductal than after intrauterine insemination (61 vs. 39%) and with high (10^6 and 10^7) rather than low (10^4 and 10^5) doses of spermatozoa for intra-uterine but not for oviductal inseminations. Researchers at our facility have achieved for the first time, the use of OI to obtain pregnancies in the mare. Fourteen mares were inseminated by OI with 50×10^3 pms

and 15 were inseminated by intrauterine AI with 500×10^6 pms. Pregnancy rates were not different between groups 3/14 (21.4%) and 6/15 (40%), respectively. Oviductal insemination has also been successfully used to obtain pregnancies in women and rabbits.

5 In cows, the site of seminal deposition during artificial insemination for the past four decades has been the uterine body. This is an acceptable technique when high numbers of fertile spermatozoa are available for insemination, but for equines -- especially when limited numbers of sperm are available -- an alternative approach has been developed. Deep intrauterine insemination is a technique that has been used to obtain pregnancies in cattle. One study compared pregnancy rates to AI when semen was deposited into the uterine body
10 or into both uterine horns (cornual insemination). Pregnancy rates when semen was deposited into the uterine body were 44.7% compared to 64.6% with cornual insemination. However, not all studies show an advantage with this technique of insemination.

As this invention shows, there can be a congruence of methods of sexing sperm based on DNA content, high speed flow cytometer/cell sorters, and procedures for inseminating
15 equines with fewer than twenty-five million total sperm without compromising fertility which may result in the possibility of a viable non-surgical or even sexed semen industry in equines. Interestingly, rather than inseminating within the uterine body where such insemination are usually placed, by insemination deep within the mare's uterine horn, better results may be achieved. By deep, it should be understood that the insertion is placed well into the uterine
20 horn. It may, but does not need to be done using the embryo transfer equipment.

As a result of the insemination, it is of course desired that an animal of the desired sex be produced. This animal may be produced according to the systems discussed earlier through the use of the sexed sperm specimen. It should also be understood that the techniques of the present invention may find application in other techniques such as
25 laproscopic insemination, oviductal insemination, or the like. As examples, the following experiments have been conducted. While not all use every aspect of the inventions described here, and do not show all the performance enhancements of the invention, they do show some enhancements possible through differing aspects of the invention.

Mares - Sixty-one reproductively normal cycling mares of light horse breeds, ranging in age from 3 to 15 were used. Mares were administered cloprostenol (250 μ l i.m.) to induce luteolysis and examined by palpation and ultrasonography of the reproductive tract *per rectum*, every other day until a follicle > 30 mm was detected, and then daily until ovulation.

- 5 Once a mare developed a follicle ≥ 35 mm, a gonadotropin releasing hormone (GnRH) implant (deslorelin acetate 2.2 mg, Ovuplant®, Fort Dodge, IA) was administered subcutaneously, and she was assigned to 1 of 3 treatment groups.

- 10 Treatment Group 1 - Mares were inseminated on a single occasion with 500×10^6 pms in a volume of 20 ml (25×10^6 pms/ml), either 40 hr (n=9) or 34 hr (n=11) after GnRH administration. Semen was deposited into the uterine body using a flexible plastic artificial insemination (AI) pipette (IMV, France).

- 15 Treatment Group 2 - Mares were inseminated on a single occasion with 25×10^6 pms in a volume of 1 ml (25×10^6 pms/ml), either 40 hr (n=13) or 34 hr (n=8) after GnRH administration. Semen was deposited at the tip of the uterine horn, ipsilateral to the preovulatory follicle, using a flexible plastic AI pipette. The location of pipette within the uterus was confirmed by transrectal ultrasonography prior to semen deposition.

- 20 Treatment Group 3 - Mares were inseminated on a single occasion with 5×10^6 pms in a volume of either 1 ml (5×10^6 pms/ml), 40 hr (n=10) or 0.2 ml (25×10^6 pms/ml), 34 hr (n=10) after GnRH administration. Mares receiving 1 ml were inseminated with a flexible plastic AI pipette, while mares receiving 0.2 ml were inseminated using a disposable implant gun (Veterinary Concepts, Green Valley, WI) containing a 0.5 ml plastic straw. Different insemination pipettes were used to optimize delivery of the two different volumes. Semen was deposited at the tip of the uterine horn, ipsilateral to the preovulatory follicle. The location of pipettes within the uterus was confirmed by transrectal ultrasonography prior to semen deposition.
- 25

After insemination, mares were examined daily to determine the day of ovulation. Pregnancy exams were performed by ultrasonography on days 12, 14, and 16 post-ovulation.

Pregnancy rates were not different between two arabian breeding stallions (Stallion A = 22/31, 71 %; Stallion B = 15/30, 50%) ($P>0.1$), or between mares bred 34 vs. 40 hours after GnRH administration (19/29, 65% and 18/32, 56%, respectively) ($P>0.1$), so the data sets were combined. As shown in Table 1, mares bred with 500×10^6 pms in a 20 ml volume had a significantly higher ($P<0.05$) pregnancy rate than mares bred with 25 or 5×10^6 pms (Table 1). There was no significant difference ($P>0.05$) in pregnancy rates between mares bred with 25×10^6 pms and mares bred with 5×10^6 pms in a volume of 1 or 0.2 mls. Although fertility was significantly higher with 500×10^6 pms when compared to Group 2 (25×10^6 pms), an initial rate of 57% was achieved with a single insemination. This was not different than pregnancy rates achieved with Group 3 (5×10^6 pms), 7/20 (35%).

Table 1 Pregnancy Rates from a Single Insemination

No. Progressively Motile Sperm	% Pregnant at Day 16
500×10^6 in 20 ml	18/20 (90%) ^a
25×10^6 in 1 ml	12/21 (57%) ^b
5×10^6 in 1 ml	3/10 (30%) ^b
5×10^6 in 0.2 ml	4/10 (40%) ^b

Values with different superscripts differ ($P<0.05$), (Chi square)

Table 1

The timing of insemination relative to GnRH administration was changed from 40 to 34 hr post GnRH during the experiment because many mares were ovulating prior to planned insemination, and therefore were not inseminated. Data from 22 mare cycles (26.5%) were excluded because they either ovulated prior to planned insemination ($n=11$), did not ovulate ($n=3$), or ovulated > 4 days after GnRH administration ($n=8$).

The optimal number of sperm generally recommended per insemination dose is 500×10^6 pms every other day while the mare is in estrus. However, as mentioned earlier, there have been studies that have shown no decrease in fertility when inseminating with 100×10^6 compared to 500×10^6 motile sperm. Studies with 50×10^6 motile sperm have shown a decrease in fertility when compared to 100 and 500×10^6 . Other studies have shown that as the number of inseminations increases, fertility increases. Unfortunately, results from these studies have been somewhat inconsistent. Therefore the present invention is directed toward

achieving the lowest number of sperm required to provide reasonable pregnancy rates when administered on a single occasion, close to ovulation.

5 In Group 3, 20 mares were inseminated with 5×10^6 pms in a volume of either 1 ml (5×10^6 pms/ml) (n= 10), or a volume of 0.2 ml (25×10^6 pms/ml) (n=10). Pregnancy rates between the two sub-groups were compared because pregnancy rates have been reported to decrease when diluting semen to a sperm concentration of $< 25 \times 10^6$ /ml. However, in this experiment there was no difference in fertility between the two sperm concentrations.

10 If a mare had already ovulated based on rectal palpation and ultrasound examination the morning of the day of planned insemination, she was not inseminated. Instead, cloprostenol (250 μ g) was administered 5 days post ovulation to induce luteolysis so she could be reused. Pregnancies were terminated at day-16 by locating by transrectal ultrasonography and disruption of the embryonic vesicle. Cloprostenol (250 μ g, i.m.) was then administered to induce luteolysis so they could be reused.

15 Semen was deposited at the tip of the uterine horn for the two lower doses in this experiment. Seminal deposition deep into the uterine horn is particularly useful when using low sperm numbers in a low volume. The flexible insemination pipette was placed in the uterus per vagina and then slowly guided to the tip of the desired uterine horn by gentle manipulation *per rectum*. The location of the pipette was confirmed by transrectal palpation and ultrasound examination.

20 An additional small, study was conducted at the end of the breeding season using five mares and one of the same two stallions. The objective of the study was to determine pregnancy rates with 25×10^6 pms deposited in the uterine body. Three of five mares (60%) inseminated on a single occasion with 25×10^6 pms 40 hr post GnRH administration were pregnant at 16 d.

In summary, the results of these experiments showed that a day-16 pregnancy rate of 57% was achieved with a single insemination, close to ovulation, With 25×10^6 pms when deposited deep into the uterine horn.

5 The objectives of the following experiments were to 1) determine pregnancy rates following insemination with 25×10^6 live-sorted, sexed spermatozoa deposited at the tip of the uterine horn ipsilateral to the preovulatory follicle and 2) compare pregnancy rates for semen sorted into a skim milk extender with or without egg-yolk.

10 Mares - Seventeen reproductively normal cycling mares of light horse breeds, ranging from 5 to 12 years of age were used. Mares were administered cloprostenol ($250 \mu\text{g i.m.}$) to induce luteolysis and examined by palpation and ultrasonography of the reproductive tract *per rectum* every other day until a follicle > 30 mm was detected, and then every day until ovulation. Once a mare developed a follicle ≥ 35 mm, she was administered a gonadotropin releasing hormone (GnRH) implant (deslorelin acetate 2.2 mg, Ovuplant®, Fort Dodge, IA) subcutaneously, and randomly assigned to 1 of 2 treatment groups.

15 Treatment Group A - Mares ($n=11$) were inseminated on a single occasion with $\sim 25 \times 10^6$ live-sorted spermatozoa in a volume of 1 ml (25 million/ml), 34 hr after GnRH administration. Spermatozoa were sorted into a commercial skim milk semen extender (EZ-Mixin, OF, Animal Reproduction Systems, Chino, CA), and the same extender was added after centrifugation as a post-centrifuge buffer to adjust sperm concentration to 25×10^6 ml.

20 Sperm were deposited at the tip of the uterine horn, ipsilateral to the preovulatory follicle, using a flexible plastic AI pipette (IMV, France). The location of pipette within the uterus was confirmed by transrectal ultrasonography prior to semen deposition. One mare was inseminated with 20×10^6 live-sorted spermatozoa because of time constraints with the flow cytometer. One mare failed to ovulate and was excluded from the study.

25 Treatment Group B - Mares ($n=10$) were inseminated on a single occasion with $\sim 25 \times 10^6$ live-sorted spermatozoa in a volume of 1 ml (25 million/ml), 34 hr after GnRH administration. One mare was inseminated with 20×10^6 live-sorted spermatozoa because of

time constraints with the flow cytometer. Spermatozoa were sorted into the same commercial semen extender plus 4% egg-yolk, and the same extender was added after centrifugation as a post-centrifuge buffer to adjust sperm concentration. Sperm were deposited at the tip of the uterine horn, ipsilateral to the preovulatory follicle using a flexible plastic AI pipette. The location of pipette within the uterus was confirmed by transrectal ultrasonography prior to semen deposition.

After insemination, mares were examined on a daily basis to determine the day of ovulation. Pregnancy exams were performed by ultrasonography on days 12, 14, 16, and 30 and post-ovulation, and fetuses were sexed on day 60.

Semen Collection and Preparation - Two stallions of Arabian breeding and known, high fertility were used in this experiment, one of which (Stallion A) was used in Experiment 1. Semen was collected the morning of planned insemination with a Colorado model artificial vagina (Animal Reproduction Systems, Chino, CA) equipped with an in-line gel filter. Ejaculates were evaluated for gel free-volume, motility and spermatozoal concentration. Semen was extended 1:1 in HBGM-3 with BSA and within minutes, transported at ambient temperature to the laboratory for further processing. The semen was centrifuged for 10 minutes at 400 x g at 22°C to highly concentrate the sperm. After centrifugation, the supernatant was aspirated, leaving a soft sperm pellet. The concentration of spermatozoa was determined using a densimeter (Animal Reproduction Systems, Chino, CA) and spermatozoa were subsequently diluted to 400 x 10⁶/ml in HBGM-3 in a total volume of 1 ml, and stained with 25 µl Hoechst 33342 (5 mg/ml water). A total of eight sample tubes were prepared and incubated at 34°C for 1 hour. Next, the stained samples were diluted to 100 x 10⁶/ml with 3 ml of HBGM-3. Food coloring (2 µl/ml of 1% FD&C #40 in HBGM-3) was added to each of the eight sample tubes, resulting in a 4 ml total volume. The samples were then filtered through a 1ml, 40 micron filter apparatus into 6 ml polypropylene tubes and held at ambient temperature until approximately 25 x 10⁶ live spermatozoa were sorted for DNA by flow cytometry. An argon laser, emitting 150 mW at 351 and 364 nm, was used on each of two MoFlo® flow cytometer/cell sorters modified for sperm sorting, operating at 50 psi with HBGM-3 without BSA as sheath fluid. Spermatozoa were collected at approximately 900 live

sperm/sec into a total of 6 polypropylene tubes (14 mls each) which contained 4 ml catch fluid before the start of sorting of either EZ-Mixin® or 4% egg- yolk in EZ-Mixin®. When two mares were available for insemination on the same day, both X- and Y-chromosome enriched sperm were collected. Tube contents were mixed every 30 minutes during sorting.

5 After sorting, sperm were pooled together from the two flow cytometers, placed in 50 ml centrifuge tubes and centrifuged for 20 minutes at 1200 x g at 22°C. The supernatant was then aspirated down to a 200µl sperm pellet, and 100µl of post-centrifuge buffer of either EZ-Mixin® CST (Animal Reproduction Systems, Chino CA) or 4% skim milk-egg yolk was added to the pellet and transferred to a 50 ml preweighed Falcon tube. A hemacytometer
10 count was done to determine final sperm concentration/ml. The volume of sperm in the Falcon tube x sperm concentration/ml equalled the total number of sperm recovered. Samples were then diluted to a total of 25×10^6 live sorted spermatozoa in a volume of 1 ml which was used for insemination.

Reanalysis of Sperm for DNA content - The relative DNA content of the sorted intact sperm
15 used for insemination was determined by flow cytometric analysis of sperm nuclei from a sample containing < 0.5 ml of each of the respective batches collected at the end of the day. Sperm nuclei were prepared from an aliquot of intact sorted sperm by sonication for 3 seconds with an Ultrasonic Dismembrator 60 (Fisher Scientific) set at setting # 2 (approximately 1 watt). The proportion of X- and Y-bearing sperm was determined by fitting
20 a pair of Gaussian distributions to the histograms from the 0° detector (Johnson et al., 1987b). Reanalysis for DNA indicated an average sort purity of 90% for X and 84% for Y chromosomebearing sperm for the 17 sorts.

Fetal Sex Determination - Fetuses from mares pregnant 60-70 days post-ovulation were sexed via transrectal ultrasonography without knowledge of the sex of the sorted sperm
25 inseminated. A real-time ultrasound scanner (Aloka 500®) equipped with a linear-array 5-Mhz transducer was used for sex determination. Fetal gender can be accurately (up to 99%) determined in horses and cattle by identifying and locating the genital tubercle (Curran, 1998).

Statistical Analysis - Data were analyzed using Fishers Exact Test

Pregnancy rates at day 16 are shown in Table 2. Pregnancy rates were not different between stallions (Stallion A = 3/10, 30%; Stallion B = 5/10, 50%) ($P > 0.1$), so the data sets were combined. There was no statistical difference in pregnancy rates between sperm treatments (EZ-Mixin = 3/10, 30% vs. 4% EY + EZ-Mixin = 5/10, 50%) ($P > 0.1$) although this result may not always be true. The phenotypic sex ratio was predicted with perfect accuracy, five out of five.

Three mares lost their pregnancy sometime between 16-60 days post-ovulation, so fetal sex could not be determined. One mare inseminated with X- bearing spermatozoa was euthanized at day 66 of gestation due to a gastro-intestinal problem. A phenotypically normal female fetus (the correct sex) was detected at necropsy.

Table 2 Pregnancy Rates Following Insemination With 25×10^6 Sexed Spermatozoa

Treatment Group	No. Mares Inseminated	No. Mares Pregnant at 16 d	No. Mares Pregnant at 60 d	Predicted*		Actual	
				% ♂	% ♀	♂	♀
EZ-Mixin	10	3 ^a	1	78	89	**	1/1
4% EY + EZ-Mixin	10	5 ^a	4	84	87	3/3	

1/1

^a No significant difference ($P > 0.1$).

*Results of reanalysis for relative DNA content of aliquots of sorted X- and Y-bearing sperm populations.

** Lost pregnancy prior to sex determination

Table 2

RESULTS: Many attempts have been made during the past 80 years to separate X- and Y-chromosome bearing sperm. The only non-destructive method that has a proven record of accurately identifying X and Y chromosome-bearing sperm is flow cytometry/cell sorting, thus making it possible to alter the sex ratio as desired. Sperm have been separated by flow cytometry/cell sorting to obtain pregnancy following surgical insemination in the following species: rabbits, swine, and horses. Surgical insemination was chosen in these experiments because of the necessity for minimizing sperm numbers due to the slow flow sorting rate

(~100 sperm/sec) of X- and Y-bearing sperm and the apparent need for large numbers of sperm to establish a pregnancy. Production of X- and Y-bearing sperm per unit time by means of high speed sorting and a newly developed orienting nozzle has increased sorting rates to 10-12 times that of previous rates. This technology has increased the number of sorted sperm per unit time and has enabled researchers to obtain pregnancies resulting from non-surgical, intra-uterine insemination in sheep and cattle.

The present study was the first to obtain viable pregnancies in the horse following non-surgical, intra-uterine insemination with sexed semen. The pregnancy rate at day 16 following insemination of 25×10^6 sexed spermatozoa (40%), was not statistically different ($P > 0.1$) than that of mares in Experiment 1 inseminated with 25×10^6 non-sorted, progressively motile spermatozoa (57%). The insemination technique was the same in both experiments. The same mares and technicians were used in both experiments. Also, both experiments were conducted during the same breeding season, at the same time of year.

Initial experimental pregnancy rates were slightly lower with sexed semen probably because of the amount of time it takes to sort 25×10^6 sperm and possible damage to the sperm by the process. In the experiments, the average time from semen collection to insemination was 7 hours. In the first experiment, mares were inseminated almost immediately after semen collection. The average total and progressive motility for the sexed spermatozoa was 69 and 38% respectively, and a total of only 25×10^6 live-sorted sperm cells were collected for insemination. The sorting process is a very stressful procedure to sperm. Sperm are pumped through fine tubing at high pressure which causes them to exit at ~100 km/hr, and stored at ambient temperature for hours until adequate numbers of sperm are collected. Sperm are incubated for one hour at 35°C with Hoechst 33342, which has a high affinity for AT-rich regions of DNA and then exposed to ultraviolet laser light at 351 and 364 nm. Unlike many DNA-specific stains, Hoechst 33342 does not intercalate into the DNA helix. While none of these processes is conducive to sperm health, no increased incidence of genetic abnormalities has been reported in the hundreds of offspring that have been produced utilizing this technology.

Of ancillary interest is the fact that others have proposed another possible explanation for lower pregnancy rates with sexed semen. They found that the first cell cycle was delayed in rabbit embryos fertilized by sperm treated with Hoechst 33342. Unfortunately, the mechanism is not known, but could be due to interference of the dye molecules as DNA is replicated or transcribed. Decreased embryo survival also has been documented in flow-sorted sperm.

Three of eight mares (38%) inseminated with sexed semen lost their pregnancies between 16-60 days. Two of these mares developed embryonic vesicles which appeared normal until day 16. The vesicles then decreased in size until they were no longer present. One of the three mares developed a viable pregnancy with a visible fetus Adth a heartbeat. The fetus was observed to be alive at day 35, but was lost by day 50. With fresh, non-sorted semen, early embryonic loss has been found to be 9% by day 14 and up to 16% on average between days 20 and 50. A sperm staining and sorting procedure was used in the present experiments. It is possible that equine sperm are more sensitive to the staining and sorting procedures than bovine sperm.

In summary, this invention has demonstrated for the first time, that pregnancy in the mare can be achieved, and foals of predetermined sex can be obtained, by deposition of a low number of spermatozoa at the tip of the uterine horn of the mare. Sexing mammalian sperm is moving away from a research technique and may now be available for commercial equine AI programs. Further, as mentioned and as can be seen from the various experiments, the field is statistically based and thus a variety of additional experiments may be conducted to further evidence the appropriate combination and limitation strategies.

The discussion included in this application is intended to serve as a basic description. The reader should be aware that the specific discussion may not explicitly describe all embodiments possible; many alternatives are implicit. It also may not fully explain the generic nature of the invention and may not explicitly show how each feature or element can actually be representative of a broader function or of a great variety of alternative or equivalent elements. Again, these are implicitly included in this disclosure. Where the

invention is described in device-oriented terminology, each element of the device implicitly performs a function. Apparatus claims may not only be included for the device described, but also method or process claims may be included to address the functions the invention and each element performs. Neither the description nor the terminology is intended to limit the scope of the claims which may be submitted. It should be understood that a variety of changes may be made without departing from the essence of the invention. Such changes are also implicitly included in the description. They still fall within the scope of this invention. A broad disclosure encompassing both the explicit embodiment(s) shown, the great variety of implicit alternative embodiments, and the broad methods or processes and the like are encompassed by this disclosure.

In addition, each of the various elements of the invention and claims may also be achieved in a variety of manners. This disclosure should be understood to encompass each such variation, be it a variation of an embodiment of any apparatus embodiment, a method or process embodiment, or even merely a variation of any element of these. Particularly, it should be understood that as the disclosure relates to elements of the invention, the words for each element may be expressed by equivalent apparatus terms or method terms -- even if only the function or result is the same. Such equivalent, broader, or even more generic terms should be considered to be encompassed in the description of each element or action. Such terms can be substituted where desired to make explicit the implicitly broad coverage to which this invention is entitled. As but one example, it should be understood that all actions may be expressed as a means for taking that action or as an element which causes that action. Similarly, each physical element disclosed should be understood to encompass a disclosure of the action which that physical element facilitates. As but one example of this aspect, the disclosure of a "collector" should be understood to encompass disclosure of the act of "collecting" -- whether explicitly discussed or not -- and, conversely, were there only disclosure of the act of "collecting", such a disclosure should be understood to encompass disclosure of a "collector." Such changes and alternative terms are to be understood to be explicitly included in the description. Further, it should be understood that in addition to the claims initially presented, the claims may be varied to more expansively address variations of each of these devices and methods set forth, each feature, component, and step shown as

separate and independent inventions, and the various combinations and permutations of each of the above.

Finally, throughout this specification -- especially the claims -- unless the context requires otherwise, the word "comprise" or variations such as "comprises" or "comprising", will be understood to imply the inclusion of a stated element or group of elements but not the exclusion of any other element or group of elements.

5

VI. CLAIMS

What is claimed is:

- 1 A method of practically producing an equine mammal comprising the steps of:
- 5 a. determining an estimated time of estrus of a female species of an equine mammal, said female species having two uterine horns, each uterine horns having a tip and a follicle, and having a vagina, a uterus, and a rectum;
- b. collecting equine sperm cells from a male species of an equine mammal;
- c. establishing an equine insemination sample containing at least some of said equine sperm cells from said male species of said equine mammal;
- 10 d. establishing a flexible probe having a sperm container;
- e. placing said flexible probe in the vagina of said female equine mammal;
- f. manipulating said flexible probe into said uterus of said female equine mammal;
- g. guiding said flexible probe into a uterine horn of said female equine mammal;
- 15 and
- h. gently manipulating said flexible probe per rectum as it is guided deep within said uterine horn of said female equine mammal to a location deep within said uterine horn of said female species near the tip of said uterine horn;
- i. artificially inseminating said female species of said equine mammal;
- 20 j. fertilizing at least one equine egg within said female species of said equine mammal; and
- k. producing an equine offspring mammal.
- 2 A method of practically producing an equine mammal as described in claim 1 wherein said step of establishing an equine insemination sample containing at least some of
- 25 said equine sperm cells from said male species of said equine mammal comprises the steps of:
- a. determining the sex characteristic of a plurality of said equine sperm cells; and
- b. sorting said equine sperm cells according to the determination of their sex characteristic,

and wherein said step of producing an equine offspring mammal comprises the step of producing an equine offspring mammal of the desired sex.

3 A method of practically producing an equine mammal as described in claim 2 wherein
5 said step of establishing an equine insemination sample containing at least some of
said equine sperm cells from said male species of said equine mammal comprises the
step of establishing an equine insemination sample containing at least some of said
equine sperm cells from said male species of said equine mammal and having a low
number of said equine sperm cells relative to the typical artificial insemination
dosage.

10 4 A method of practically producing an equine mammal as described in claim 3 wherein
said step of establishing an equine insemination sample containing at least some of
said equine sperm cells from said male species of said equine mammal and having a
low number of said equine sperm cells relative to the typical artificial insemination
15 dosage comprises the step of establishing an equine insemination sample containing at
least some of said equine sperm cells from said male species of said equine mammal
and having a low number of said equine sperm cells relative to the typical artificial
insemination dosage selected from the group consisting of: an equine insemination
sample of no more than about five million sperm cells, and an equine insemination
sample of no more than about twenty-five million sperm cells.

20 5 A method of practically producing an equine mammal as described in claim 4 wherein
said step of fertilizing at least one equine egg within said female species of said
equine mammal comprises the step of fertilizing at least one equine egg within said
female species of said equine mammal at success levels statistically comparable to the
typical artificial insemination dosage.

25 6 A method of practically producing an equine mammal as described in claim 5 wherein
said step of establishing an equine insemination sample containing at least some of
said equine sperm cells from said male species of said equine mammal and having a

low number of said equine sperm cells relative to the typical artificial insemination dosage comprises the steps of:

- a. staining said equine sperm cells;
- b. sorting according to said sex of said equine sperm cells through the use of high speed flow cytometry; and
- c. concentrating said sorted equine sperm cells.

7 A method of practically producing an equine mammal as described in claim 6 wherein said step of sorting according to said sex of said equine sperm cells through the use of high speed flow cytometry comprises the step of collecting live sperm of the desired sex at the rate of at least nine hundred live sperm per second.

8 A method of practically producing an equine mammal as described in claim 6 wherein said step of sorting according to said sex of said equine sperm cells through the use of high speed flow cytometry comprises the step of operating a high speed cell sorter at a pressure of at least about fifty pounds per square inch.

15 9 A method of practically producing an equine mammal as described in claim 1 wherein said step of establishing an equine insemination sample containing at least some of said equine sperm cells from said male species of said equine mammal comprises the step of establishing an equine insemination sample containing at least some of said equine sperm cells from said male species of said equine mammal and having a low number of said equine sperm cells relative to the typical artificial insemination dosage.

10 25 A method of practically producing an equine mammal as described in claim 9 wherein said step of establishing an equine insemination sample containing at least some of said equine sperm cells from said male species of said equine mammal and having a low number of said equine sperm cells relative to the typical artificial insemination dosage comprises the step of establishing an equine insemination sample containing at least some of said equine sperm cells from said male species of said equine mammal

and having a low number of said equine sperm cells relative to the typical artificial insemination dosage selected from the group consisting of: an equine insemination sample of no more than about five million sperm cells, and an equine insemination sample of no more than about twenty-five million sperm cells.

- 5 11 A method of practically producing an equine mammal as described in claim 4 wherein said step of establishing an equine insemination sample containing at least some of said equine sperm cells from said male species of said equine mammal comprises the step of establishing an equine insemination sample having a volume selected from the group: 0.2 ml or 1ml.
- 10 12 A method of practically producing an equine mammal as described in claim 7 wherein said step of establishing an equine insemination sample containing at least some of said equine sperm cells from said male species of said equine mammal comprises the step of establishing an equine insemination sample having a volume selected from the group: 0.2 ml, or 1ml.
- 15 13 A method of practically producing an equine mammal as described in claim 10 wherein said step of establishing an equine insemination sample containing at least some of said equine sperm cells from said male species of said equine mammal comprises the step of establishing an equine insemination sample having a volume selected from the group: 0.2 ml, or 1ml.
- 20 14 An animal produced through use of a method as described in any of claims 1, 4, 7, 11, or 12.
- 15 15 A method of practically producing an equine mammal comprising the steps of:
- 25 a. determining an estimated time of estrus of a female species of an equine mammal, said female species having two uterine horns, each uterine horns having a tip and a follicle;
- b. collecting equine sperm cells from a male species of an equine mammal;

- c. establishing an equine insemination sample containing at least some of said equine sperm cells from said male species of said equine mammal;
- d. inserting at least a portion of said equine insemination sample deep within at least one of said uterine horns of said female species near the tip of said uterine horn;
- e. artificially inseminating said female species of said equine mammal;
- f. fertilizing at least one equine egg within said female species of said equine mammal; and
- g. producing an equine offspring mammal.

5

10

16

A method of practically producing an equine mammal as described in claim 15 and further comprising the step of ascertaining which uterine horn is ipsilateral to the preovulatory follicle and wherein said step of inserting at least a portion of said equine insemination sample deep within at least one of said uterine horns of said female equine species comprises the step of inserting at least a portion of said equine insemination sample near the tip of said uterine horn ipsilateral to the preovulatory follicle.

15

17

A method of practically producing an equine mammal as described in claim 15 wherein said step of artificially inseminating said female species of said equine mammal comprises the step of artificially inseminating said female equine species on a single occasion close to ovulation.

20

18

A method of practically producing an equine mammal as described in claim 17 wherein said step of artificially inseminating said female equine species on a single occasion close to ovulation comprises the step of artificially inseminating said female equine species both ipsi- and contra-lateral within the uterine horns of said equine mammal.

25

19

A method of practically producing an equine mammal as described in claim 15 wherein said steps of inserting at least a portion of said equine insemination sample

deep within at least one of said uterine horns of said female species near the tip of said uterine horn, artificially inseminating said female species of said equine mammal, and fertilizing at least one equine egg within said female species of said equine mammal are each accomplished in a field environment.

- 5 20 A method of practically producing an equine mammal as described in claim 15 wherein said step of inserting at least a portion of said equine insemination sample deep within at least one of said uterine horns of said female species near the tip of said uterine horn comprises the step of inserting at least a portion of said equine insemination sample at a location selected from the group consisting of: within about one-half inch of the tip of said uterine horn, within about one inch of the tip of said uterine horn, and within about two inches of the tip of said uterine horn.
- 10
- 21 A method of practically producing an equine mammal as described in claim 15 and further comprising the step of manipulating the ovulation of said female equine mammal prior to accomplishing the step of artificially inseminating said female species of said equine mammal.
- 15
- 22 A method of practically producing an equine mammal as described in claim 21 wherein said step of manipulating the ovulation of said female equine mammal prior to accomplishing the step of artificially inseminating said female species of said equine mammal comprises the step of administering a gonadotropin releasing hormone to said female equine mammal.
- 20
- 23 A method of practically producing an equine mammal as described in claim 22 wherein said step of artificially inseminating said female species of said equine mammal is accomplished at a time selected from the group consisting of: about thirty four hours after said step of administering said gonadotropin releasing hormone to said female equine mammal, about forty hours after said step of administering said gonadotropin releasing hormone to said female equine mammal, and between about
- 25

thirty four hours to about forty hours after said step of administering said gonadotropin releasing hormone to said female equine mammal.

24 A method of practically producing an equine mammal as described in claim 21
5 wherein said step of inserting at least a portion of said equine insemination sample deep within at least one of said uterine horns of said female species near the tip of said uterine horn comprises the step of inserting said equine insemination sample within said uterine horn through the use of a flexible plastic artificial insemination pipette.

25 A method of practically producing an equine mammal as described in claim 15
10 wherein said step of inserting at least a portion of said equine insemination sample deep within at least one of said uterine horns of said female species near the tip of said uterine horn comprises the steps of:

- a. establishing a flexible probe having a sperm container;
- b. placing said flexible probe in the vagina of said female equine mammal;
- c. manipulating said flexible probe into said uterus of said female equine
15 mammal;
- d. slowly guiding said flexible probe into a uterine horn of said female equine mammal; and
- e. gently manipulating said flexible probe per rectum as it is guided deep within
20 said uterine horn of said female equine mammal to a location deep within said uterine horn of said female species near the tip of said uterine horn.

26 A method of practically producing an equine mammal as described in claim 15 and
further comprising the step of confirming by transrectal ultrasonography the location
at which said step of inserting at least a portion of said equine insemination sample is
accomplished prior to accomplishing said step of artificially inseminating said female
25 species of said equine mammal.

27 A method of practically producing an equine mammal as described in claim 15
wherein said step of establishing an equine insemination sample containing at least

5 some of said sperm cells from said male species of said equine mammal comprises the step of establishing an equine insemination sample having a low number of said equine sperm cells relative to the typical artificial insemination dosage, and wherein said step of fertilizing at least one equine egg within said female species of said equine mammal comprises the step of fertilizing at least one equine egg within said female species of said equine mammal at success levels statistically comparable to the typical artificial insemination dosage.

10 28 A method of practically producing an equine mammal as described in claim 27 wherein said step of establishing an equine insemination sample having a low number of said equine sperm cells relative to the typical artificial insemination dosage comprises the step of establishing an equine insemination sample selected from the group consisting of: an equine insemination sample of no more than about five million sperm cells, and an equine insemination sample of no more than about twenty-five million sperm cells.

15 29 A method of practically producing an equine mammal as described in claim 25 wherein said step of establishing an equine insemination sample containing at least some of said sperm cells from said male species of said equine mammal comprises the step of establishing an equine insemination sample having a low number of said equine sperm cells relative to the typical artificial insemination dosage, and wherein
20 said step of fertilizing at least one equine egg within said female species of said equine mammal comprises the step of fertilizing at least one equine egg within said female species of said equine mammal at success levels statistically comparable to the typical artificial insemination dosage.

25 30 A method of practically producing an equine mammal as described in claim 29 wherein said step of establishing an equine insemination sample having a low number of said equine sperm cells relative to the typical artificial insemination dosage comprises the step of establishing an equine insemination sample selected from the group consisting of: an equine insemination sample of no more than about five million

sperm cells, and an equine insemination sample of no more than about twenty-five million sperm cells.

31 A method of practically producing an equine mammal as described in claim 29 wherein said step of establishing an equine insemination sample containing at least some of said equine sperm cells from said male species of said equine mammal comprises the steps of:

- a. determining the sex characteristic of a plurality of said equine sperm cells; and
- b. sorting said equine sperm cells according to the determination of their sex characteristic,

and wherein said step of producing an equine offspring mammal comprises the step of producing an equine offspring mammal of the desired sex.

32 A method of practically producing an equine mammal as described in claim 31 wherein said steps of determining the sex characteristic of a plurality of said equine sperm cells and sorting said equine sperm cells according to the determination of their sex characteristic comprise the steps of:

- a. staining said equine sperm cells;
- b. sorting according to said sex of said equine sperm cells through the use of high speed flow cytometry; and
- c. concentrating said sorted equine sperm cells.

33 A method of practically producing an equine mammal as described in claim 32 wherein said step of sorting according to said sex of said equine sperm cells through the use of high speed flow cytometry comprises the step of collecting live sperm of the desired sex at the rate of at least nine hundred live sperm per second.

34 A method of practically producing an equine mammal as described in claim 32 wherein said step of sorting according to said sex of said equine sperm cells through the use of high speed flow cytometry comprises the step of operating a high speed cell sorter at a pressure of at least about fifty pounds per square inch.

35 A method of practically producing an equine mammal as described in claim 32 wherein said step of sorting according to said sex of said equine sperm cells through the use of high speed flow cytometry comprises the step of collecting equine sperm cells having the desired sex characteristic in a skim milk solution.

5 36 A method of practically producing an equine mammal as described in claim 33 wherein said step of establishing an equine insemination sample containing at least some of said sperm cells from said male species of said equine mammal comprises the step of establishing an equine insemination sample having a low number of said equine sperm cells relative to the typical artificial insemination dosage, and wherein
10 said step of fertilizing at least one equine egg within said female species of said equine mammal comprises the step of fertilizing at least one equine egg within said female species of said equine mammal at success levels statistically comparable to the typical artificial insemination dosage.

15 37 A method of practically producing an equine mammal as described in claim 36 wherein said step of establishing an equine insemination sample having a low number of said equine sperm cells relative to the typical artificial insemination dosage comprises the step of establishing an equine insemination sample selected from the group consisting of: an equine insemination sample of no more than about five million sperm cells, and an equine insemination sample of no more than about twenty-five
20 million sperm cells.

38 A method of practically producing an equine mammal as described in claim 15 wherein said step of establishing an equine insemination sample containing at least some of said equine sperm cells from said male species of said equine mammal comprises the steps of:
25 a. determining the sex characteristic of a plurality of said equine sperm cells; and
b. sorting said equine sperm cells according to the determination of their sex characteristic,

and wherein said step of producing an equine offspring mammal comprises the step of producing an equine offspring mammal of the desired sex.

39 A method of practically producing an equine mammal as described in claim 25 wherein said step of establishing an equine insemination sample containing at least some of said equine sperm cells from said male species of said equine mammal comprises the steps of:

- a. determining the sex characteristic of a plurality of said equine sperm cells; and
- b. sorting said equine sperm cells according to the determination of their sex characteristic,

and wherein said step of producing an equine offspring mammal comprises the step of producing an equine offspring mammal of the desired sex.

40 A method of practically producing an equine mammal as described in claim 15 and further comprising the step of administering an equine pituitary extract to said female equine mammal to enhance the probability at which said step of fertilizing at least one equine egg within said female species of said equine mammal occurs.

41 A method of practically producing an equine mammal as described in claim 28 wherein said step of establishing an equine insemination sample containing at least some of said equine sperm cells from said male species of said equine mammal comprises the step of establishing an equine insemination sample having a volume selected from the group: 0.2 ml, or 1 ml.

42 A method of practically producing an equine mammal as described in claim 30 wherein said step of establishing an equine insemination sample containing at least some of said equine sperm cells from said male species of said equine mammal comprises the step of establishing an equine insemination sample having a volume selected from the group: 0.2 ml, or 1 ml.

43 A method of practically producing an equine mammal as described in claim 37
wherein said step of establishing an equine insemination sample containing at least
some of said equine sperm cells from said male species of said equine mammal
comprises the step of establishing an equine insemination sample having a volume
5 selected from the group: 0.2 ml, or 1ml.

44 An animal produced through use of a method as described in any of claims 15, 19, 23,
25, 28, 30, 31, 33, 35, 40, 41, , or 43.

45 A method of sorting equine sperm cells according to a determination of their sex
characteristic comprising the steps of:

- 10 a. collecting equine sperm cells from a male species of an equine mammal;
- b. staining said equine sperm cells;
- c. establishing a cell source which supplies said equine sperm cells to be sorted;
- d. establishing a sheath fluid which is adapted to form droplets and which is
compatible with said equine sperm cells;
- 15 e. establishing a skim milk solution into which said equine sperm cells are
collected;
- f. discriminating between said equine sperm cells according to a determination
of their sex characteristic;
- g. entraining individual equine sperm cells in a droplet;
- 20 h. sorting said droplets according to said sex of the individual equine sperm cells
they contain; and
- i. collecting equine sperm cells having the desired sex characteristic in said skim
milk solution.

46 A method of sorting equine sperm cells according to a determination of their sex
characteristic as described in claim 45 wherein said step of establishing a skim milk
solution into which said equine sperm cells are collected comprises the step of
establishing a solution containing a skim milk extender as a collection fluid.
25

- 47 A method of sorting equine sperm cells according to a determination of their sex characteristic as described in claim 46 wherein said step of establishing a skim milk solution into which said equine sperm cells are collected further comprises the step of establishing a solution containing about four percent egg yolk as a collection fluid.
- 5 48 A method of sorting equine sperm cells according to a determination of their sex characteristic as described in claim 45 wherein said step of establishing a sheath fluid which is adapted to form droplets and which is compatible with said equine sperm cells comprises the step of establishing a sheath fluid containing a hepes buffered medium.
- 10 49 A method of sorting equine sperm cells according to a determination of their sex characteristic as described in claim 45 wherein said step of sorting said droplets according to said sex of the individual equine sperm cells they contain comprises the step of sorting according to said sex of said equine sperm cells through the use of high speed flow cytometry.
- 15 50 A method of sorting equine sperm cells according to a determination of their sex characteristic as described in claim 49 wherein said step of sorting according to said sex of said equine sperm cells through the use of high speed flow cytometry comprises the step of collecting live sperm of the desired sex at the rate of at least nine hundred live sperm per second.
- 20 51 A method of sorting equine sperm cells according to a determination of their sex characteristic as described in claim 49 wherein said step of sorting according to said sex of said equine sperm cells through the use of high speed flow cytometry comprises the step of operating a high speed cell sorter at a pressure of at least about fifty pounds per square inch.
- 25 52 A method of practically producing an equine mammal involving sorted equine sperm cells wherein said sorted equine sperm cells are sorted according to a determination of

their sex characteristic through a method as described in claim 45 and further comprising the steps of:

- a. determining an estimated time of estrus of a female species of an equine mammal, said female species having two uterine horns, each uterine horns having a tip and a follicle;
- b. establishing an equine insemination sample containing at least some of said sorted equine sperm cells from said male species of said equine mammal and having a low number of said equine sperm cells relative to the typical artificial insemination dosage;
- c. non-surgically inserting at least a portion of said equine insemination sample in said female species of said equine mammal;
- d. artificially inseminating said female species of said equine mammal;
- e. fertilizing at least one equine egg within said female species of said equine mammal; and
- f. producing an equine offspring mammal of the desired sex.

53 A method of practically producing an equine mammal as described in claim 52 wherein said step of establishing an equine insemination sample containing at least some of said sorted equine sperm cells from said male species of said equine mammal and having a low number of said equine sperm cells relative to the typical artificial insemination dosage comprises the step of establishing an equine insemination sample containing at least some of said equine sperm cells from said male species of said equine mammal and having a low number of said equine sperm cells relative to the typical artificial insemination dosage selected from the group consisting of: an equine insemination sample of no more than about five million sperm cells, and an equine insemination sample of no more than about twenty-five million sperm cells.

54 A method of practically producing an equine mammal as described in claim 53 and further comprising the step of ascertaining which uterine horn is ipsilateral to the preovulatory follicle and wherein said step of non-surgically inserting at least a portion of said equine insemination sample in said female species of said equine

mammal comprises the step of inserting at least a portion of said equine insemination sample near the tip of said uterine horn ipsilateral to the preovulatory follicle.

55 A method of practically producing an equine mammal as described in claim 53
wherein said step of artificially inseminating said female species of said equine
5 mammal comprises the step of artificially inseminating said female equine species on
a single occasion close to ovulation.

56 A method of practically producing an equine mammal as described in claim 55
wherein said step of artificially inseminating said female equine species on a single
10 occasion close to ovulation comprises the step of comprises the step of artificially
inseminating said female equine species both ipsi- and contra-lateral within the
uterine horns of said equine mammal.

57 A method of practically producing an equine mammal as described in claim 54
wherein said steps of inserting at least a portion of said equine insemination sample
15 near the tip of said uterine horn ipsilateral to the preovulatory follicle, artificially
inseminating said female species of said equine mammal, and fertilizing at least one
equine egg within said female species of said equine mammal are each accomplished
in a field environment.

58 A method of practically producing an equine mammal as described in claim 53
wherein said step of establishing an equine insemination sample containing at least
20 some of said equine sperm cells from said male species of said equine mammal
comprises the step of establishing an equine insemination sample having a volume
selected from the group: 0.2 ml, or 1ml.

59 An animal produced through use of a method as described in any of claims 45, 50, 53,
57, or 58.

60 A method of flow cytometry accomplished through use of a method as described in any of claims 45, 48, or 50.

61 An equine-adapted flow cytometer system for isolating desired cells comprising:

- 5 a. an equine sperm cell source which supplies cells to be analyzed by the flow cytometer;
- b. a chemically coordinated sheath fluid source which creates a sheath fluid environment for said equine sperm cells;
- c. a nozzle through which said equine sperm cells pass while subjected to said sheath fluid environment;
- 10 d. an oscillator which acts upon said sheath fluid as it passes through said nozzle;
- e. a cell sensing system which responds to said equine sperm cells;
- f. a equine sperm cell sorter discrimination system which acts to sort said equine sperm cells having a desired sex characteristic; and
- 15 g. a skim milk solution collector into which said equine sperm cells having the desired sex characteristic are placed.

62 An equine-adapted flow cytometer system for isolating desired cells as described in claim 61 wherein said skim milk solution collector comprises a solution containing a skim milk semen extender.

20 63 An equine-adapted flow cytometer system for isolating desired cells as described in claim 62 wherein said skim milk solution collector further comprises a solution containing about four percent egg yolk.

25 64 An equine-adapted flow cytometer system for isolating desired cells as described in claim 62 wherein said chemically coordinated sheath fluid source comprises a sheath fluid containing a hepes buffered medium.

65 An equine-adapted flow cytometer system for isolating desired cells as described in claim 61 wherein said equine sperm cell sorter discrimination system comprises a high speed cell sorter.

5 66 An equine-adapted flow cytometer system for isolating desired cells as described in claim 65 wherein said a high speed cell sorter collects live sperm of the desired sex at the rate of at least nine hundred live sperm per second

67 An equine-adapted flow cytometer system for isolating desired cells as described in claim 65 wherein said a high speed cell sorter operates at a pressure of at least about fifty pounds per square inch.

10 68 A sexed sperm specimen produced through use of a system as described in any of claims 61, 62, 64, 65 or 66.

69 A sexed equine animal produced through use of a system as described in any of claims 61, 62, 64, 65 or 66.

15 70 A method of practically producing an equine mammal comprising the steps of:

- a. determining an estimated time of estrus of a female species of an equine mammal, said female species having two uterine horns, each uterine horns having a tip and a follicle;
- b. collecting equine sperm cells from a male species of an equine mammal;
- c. determining the sex characteristic of a plurality of said equine sperm cells;
- 20 d. sorting said equine sperm cells according to the determination of their sex characteristic;
- e. establishing an equine insemination sample containing at least some of said sorted equine sperm cells from said male species of said equine mammal;
- f. non-surgically inserting at least a portion of said equine insemination sample in said female species of said equine mammal;
- 25 g. artificially inseminating said female species of said equine mammal;

- h. fertilizing at least one equine egg within said female species of said equine mammal; and
- i. producing an equine offspring mammal of the desired sex.

71 A method of practically producing an equine mammal as described in claim 70
5 wherein said steps of determining the sex characteristic of a plurality of said equine sperm cells and sorting said equine sperm cells according to the determination of their sex characteristic comprise the steps of:

- a. staining said equine sperm cells;
- b. sorting according to said sex of said equine sperm cells through the use of high
10 speed flow cytometry; and
- c. concentrating said sorted equine sperm cells.

72 A method of practically producing an equine mammal as described in claim 71
15 wherein said step of sorting according to said sex of said equine sperm cells through the use of high speed flow cytometry comprises the step of collecting live sperm of the desired sex at the rate of at least nine hundred live sperm per second.

73 A method of practically producing an equine mammal as described in claim 71
wherein said step of sorting according to said sex of said equine sperm cells through the use of high speed flow cytometry comprises the step of operating a high speed cell
sorter at a pressure of at least about fifty pounds per square inch.

20 74 A method of practically producing an equine mammal as described in claim 71
wherein said step of sorting according to said sex of said equine sperm cells through the use of high speed flow cytometry comprises the step of collecting equine sperm
cells having the desired sex characteristic in a skim milk solution.

25 75 A method of practically producing an equine mammal as described in claim 70
wherein said step of establishing an equine insemination sample containing at least
some of said equine sperm cells from said male species of said equine mammal

comprises the step of establishing an equine insemination sample containing at least some of said equine sperm cells from said male species of said equine mammal and having a low number of said equine sperm cells relative to the typical artificial insemination dosage selected from the group consisting of: an equine insemination sample of no more than about five million sperm cells, and an equine insemination sample of no more than about twenty-five million sperm cells.

76 A method of practically producing an equine mammal as described in claim 72 wherein said step of establishing an equine insemination sample containing at least some of said equine sperm cells from said male species of said equine mammal comprises the step of establishing an equine insemination sample containing at least some of said equine sperm cells from said male species of said equine mammal and having a low number of said equine sperm cells relative to the typical artificial insemination dosage selected from the group consisting of: an equine insemination sample of no more than about five million sperm cells, and an equine insemination sample of no more than about twenty-five million sperm cells.

77 A method of practically producing an equine mammal as described in claim 75 wherein said step of establishing an equine insemination sample containing at least some of said equine sperm cells from said male species of said equine mammal comprises the step of establishing an equine insemination sample having a volume selected from the group: 0.2 ml, or 1ml.

78 A method of practically producing an equine mammal as described in claim 76 wherein said step of establishing an equine insemination sample containing at least some of said equine sperm cells from said male species of said equine mammal comprises the step of establishing an equine insemination sample having a volume selected from the group: 0.2 ml, or 1ml.

79 A method of practically producing an equine mammal as described in claim 70 wherein said step of non-surgically inserting at least a portion of said equine

insemination sample in said female species of said equine mammal comprises the step of inserting at least a portion of said equine insemination sample deep within at least one of said uterine horns of said female species near the tip of said uterine horn.

5 80 A method of practically producing an equine mammal as described in claim 71 wherein said step of non-surgically inserting at least a portion of said equine insemination sample in said female species of said equine mammal comprises the step of inserting at least a portion of said equine insemination sample deep within at least one of said uterine horns of said female species near the tip of said uterine horn.

10 81 A method of practically producing an equine mammal as described in claim 79 wherein said step of artificially inseminating said female species of said equine mammal comprises the step of artificially inseminating said female equine species on a single occasion close to ovulation.

15 82 A method of practically producing an equine mammal as described in claim 80 wherein said step of artificially inseminating said female species of said equine mammal comprises the step of artificially inseminating said female equine species on a single occasion close to ovulation.

20 83 A method of practically producing an equine mammal as described in claim 81 and further comprising the step of manipulating the ovulation of said female equine mammal prior to accomplishing the step of artificially inseminating said female species of said equine mammal.

84 A method of practically producing an equine mammal as described in claim 82 and further comprising the step of manipulating the ovulation of said female equine mammal prior to accomplishing the step of artificially inseminating said female species of said equine mammal.

85 A method of practically producing an equine mammal as described in claim 83 wherein said step of manipulating the ovulation of said female equine mammal prior to accomplishing the step of artificially inseminating said female species of said equine mammal comprises the step of administering a gonadotropin releasing hormone to said female equine mammal.

86 A method of practically producing an equine mammal as described in claim 84 herein said step of manipulating the ovulation of said female equine mammal prior to accomplishing the step of artificially inseminating said female species of said equine mammal comprises the step of administering a gonadotropin releasing hormone to said female equine mammal.

87 A method of practically producing an equine mammal as described in claim 85 herein said step of artificially inseminating said female species of said equine mammal is accomplished at a time selected from the group consisting of: about thirty four hours after said step of administering said gonadotropin releasing hormone to said female equine mammal, about forty hours after said step of administering said gonadotropin releasing hormone to said female equine mammal, and between about thirty four hours to about forty hours after said step of administering said gonadotropin releasing hormone to said female equine mammal.

88 A method of practically producing an equine mammal as described in claim 86 wherein said step of artificially inseminating said female species of said equine mammal is accomplished at a time selected from the group consisting of: about thirty four hours after said step of administering said gonadotropin releasing hormone to said female equine mammal, about forty hours after said step of administering said gonadotropin releasing hormone to said female equine mammal, and between about thirty four hours to about forty hours after said step of administering said gonadotropin releasing hormone to said female equine mammal.

89 A method of practically producing an equine mammal as described in claim 85 wherein said step of inserting at least a portion of said equine insemination sample deep within at least one of said uterine horns of said female species near the tip of said uterine horn comprises the steps of:

- 5 a. establishing a flexible probe having a sperm container;
- b. placing said flexible probe in the vagina of said female equine mammal;
- c. manipulating said flexible probe into said uterus of said female equine mammal;
- d. guiding said flexible probe into a uterine horn of said female equine mammal;
- 10 and
- e. gently manipulating said flexible probe per rectum as it is guided deep within said uterine horn of said female equine mammal to a location deep within said uterine horn of said female species near the tip of said uterine horn.

90 A method of practically producing an equine mammal as described in claim 86 wherein said step of inserting at least a portion of said equine insemination sample deep within at least one of said uterine horns of said female species near the tip of said uterine horn comprises the steps of:

- 15 a. establishing a flexible probe having a sperm container;
- b. placing said flexible probe in the vagina of said female equine mammal;
- 20 c. manipulating said flexible probe into said uterus of said female equine mammal;
- d. guiding said flexible probe into a uterine horn of said female equine mammal;
- and
- 25 e. gently manipulating said flexible probe per rectum as it is guided deep within said uterine horn of said female equine mammal to a location deep within said uterine horn of said female species near the tip of said uterine horn.

91 An animal produced through use of a method as described in any of claims 70, 72, 74, 77, 78, 79, or 89.

92 A method of practically producing an equine mammal comprising the steps of:

- a. determining an estimated time of estrus of a female species of an equine mammal, said female species having two uterine horns, each uterine horns having a tip and a follicle;
- 5 b. collecting equine sperm cells from a male species of an equine mammal;
- c. establishing an equine insemination sample containing at least some of said equine sperm cells from said male species of said equine mammal and having a low number of said equine sperm cells relative to the typical artificial insemination dosage;
- 10 d. non-surgically inserting at least a portion of said equine insemination sample in said female species of said equine mammal;
- e. artificially inseminating said female species of said equine mammal;
- f. fertilizing at least one equine egg within said female species of said equine mammal at success levels statistically comparable to the typical artificial insemination dosage; and
- 15 g. producing an equine offspring mammal.

93 A method of practically producing an equine mammal as described in claim 92 and further comprising the step of ascertaining which uterine horn is ipsilateral to the preovulatory follicle and wherein said step of inserting at least a portion of said equine
20 insemination sample deep within at least one of said uterine horns of said female equine species comprises the step of inserting at least a portion of said equine insemination sample near the tip of said uterine horn ipsilateral to the preovulatory follicle.

94 A method of practically producing an equine mammal as described in claim 92 wherein said step of artificially inseminating said female species of said equine
25 mammal comprises the step of artificially inseminating said female equine species on a single occasion close to ovulation.

95 A method of practically producing an equine mammal as described in claim 94
wherein said step of artificially inseminating said female equine species on a single
occasion close to ovulation comprises the step of artificially
inseminating said female equine species both ipsi- and contra-lateral within the
5 uterine horns of said equine mammal.

96 A method of practically producing an equine mammal as described in claim 92
wherein said steps of inserting at least a portion of said equine insemination sample
deep within at least one of said uterine horns of said female species near the tip of said
uterine horn, artificially inseminating said female species of said equine mammal, and
10 fertilizing at least one equine egg within said female species of said equine mammal
are each accomplished in a field environment.

97 A method of practically producing an equine mammal as described in claim 92
wherein said step of fertilizing at least one equine egg within said female species of
said equine mammal at success levels statistically comparable to the typical artificial
insemination dosage comprises the step of fertilizing at least one equine egg within
15 said female species of said equine mammal at success levels selected from the group
consisting of at least 90%, at least 81%, at least 75%, at least 65%, at least 60%, at
least 57%, at least 40%, at least 35%, and at least 30%.

98 A method of practically producing an equine mammal as described in claim 92
wherein said step of establishing an equine insemination sample having a low number
20 of said equine sperm cells relative to the typical artificial insemination dosage
comprises the step of establishing an equine insemination sample selected from the
group consisting of: an equine insemination sample of no more than about forty
million sperm cells, an equine insemination sample of no more than about fifty
25 million sperm cells, and an equine insemination sample of no more than about one
hundred million sperm cells.

99 A method of practically producing an equine mammal as described in claim 94
wherein said step of establishing an equine insemination sample having a low number
of said equine sperm cells relative to the typical artificial insemination dosage
comprises the step of establishing an equine insemination sample selected from the
5 group consisting of: an equine insemination sample of no more than about forty
million sperm cells, an equine insemination sample of no more than about fifty
million sperm cells, and an equine insemination sample of no more than about one
hundred million sperm cells.

100 A method of practically producing an equine mammal as described in claim 97
10 wherein said step of establishing an equine insemination sample having a low number
of said equine sperm cells relative to the typical artificial insemination dosage
comprises the step of establishing an equine insemination sample selected from the
group consisting of: an equine insemination sample of no more than about forty
million sperm cells, an equine insemination sample of no more than about fifty
15 million sperm cells, and an equine insemination sample of no more than about one
hundred million sperm cells.

101 A method of practically producing an equine mammal as described in claim 92
wherein said step of establishing an equine insemination sample containing at least
some of said equine sperm cells from said male species of said equine mammal
20 comprises the steps of:
a. determining the sex characteristic of a plurality of said equine sperm cells; and
b. sorting said equine sperm cells according to the determination of their sex
characteristic,
and wherein said step of producing an equine offspring mammal comprises the step of
25 producing an equine offspring mammal of the desired sex.

102 A method of practically producing an equine mammal as described in claim 94
wherein said step of establishing an equine insemination sample containing at least

some of said equine sperm cells from said male species of said equine mammal comprises the steps of:

- a. determining the sex characteristic of a plurality of said equine sperm cells; and
- b. sorting said equine sperm cells according to the determination of their sex characteristic,

and wherein said step of producing an equine offspring mammal comprises the step of producing an equine offspring mammal of the desired sex.

103 A method of practically producing an equine mammal as described in claim 97 wherein said step of establishing an equine insemination sample containing at least some of said equine sperm cells from said male species of said equine mammal comprises the steps of:

- a. determining the sex characteristic of a plurality of said equine sperm cells; and
- b. sorting said equine sperm cells according to the determination of their sex characteristic,

and wherein said step of producing an equine offspring mammal comprises the step of producing an equine offspring mammal of the desired sex.

104 A method of practically producing an equine mammal as described in claim 102 wherein said steps of determining the sex characteristic of a plurality of said equine sperm cells and sorting said equine sperm cells according to the determination of their sex characteristic comprise the steps of:

- a. staining said equine sperm cells;
- b. sorting according to said sex of said equine sperm cells through the use of high speed flow cytometry; and
- c. concentrating said sorted equine sperm cells.

105 A method of practically producing an equine mammal as described in claim 104 wherein said step of sorting according to said sex of said equine sperm cells through the use of high speed flow cytometry comprises the step of collecting live sperm of the desired sex at the rate of at least nine hundred live sperm per second.

106 A method of practically producing an equine mammal as described in claim 104 wherein said step of sorting according to said sex of said equine sperm cells through the use of high speed flow cytometry comprises the step of operating a high speed cell sorter at a pressure of at least about fifty pounds per square inch.

5 107 A method of practically producing an equine mammal as described in claim 104 wherein said step of sorting according to said sex of said equine sperm cells through the use of high speed flow cytometry comprises the step of collecting equine sperm cells having the desired sex characteristic in a skim milk solution.

10 108 An animal produced through use of a method as described in any of claims 92, 96, 97, 100, or 101.

109 A method of practically producing an equine mammal comprising the steps of:

- a. determining an estimated time of estrus of a female species of an equine mammal, said female species having two uterine horns, each uterine horns having a tip and a follicle;
- 15 b. collecting equine sperm cells from a male species of an equine mammal;
- c. establishing an equine insemination sample containing at least some of said equine sperm cells from said male species of said equine mammal and having a low number of said equine sperm cells relative to the typical artificial insemination dosage selected from the group consisting of: an equine insemination sample of no more than about five million sperm cells, and an
20 equine insemination sample of no more than about twenty-five million sperm cells;
- d. non-surgically inserting at least a portion of said equine insemination sample in said female species of said equine mammal;
- 25 e. artificially inseminating said female species of said equine mammal;
- f. fertilizing at least one equine egg within said female species of said equine mammal; and
- g. producing an equine offspring mammal.

110 A method of practically producing an equine mammal as described in claim 109 and further comprising the step of ascertaining which uterine horn is ipsilateral to the preovulatory follicle and wherein said step of non-surgically inserting at least a portion of said equine insemination sample in said female species of said equine mammal comprises the step of inserting at least a portion of said equine insemination sample near the tip of said uterine horn ipsilateral to the preovulatory follicle.

111 A method of practically producing an equine mammal as described in claim 109 wherein said step of artificially inseminating said female species of said equine mammal comprises the step of artificially inseminating said female equine species on a single occasion close to ovulation.

112 A method of practically producing an equine mammal as described in claim 111 wherein said step of artificially inseminating said female equine species on a single occasion close to ovulation comprises the step of artificially inseminating said female equine species both ipsi- and contra-lateral within the uterine horns of said equine mammal.

113 A method of practically producing an equine mammal as described in claim 109 wherein said steps of non-surgically inserting at least a portion of said equine insemination sample in said female species of said equine mammal, artificially inseminating said female species of said equine mammal, and fertilizing at least one equine egg within said female species of said equine mammal are each accomplished in a field environment.

114 A method of practically producing an equine mammal as described in claim 109 wherein said step of establishing an equine insemination sample containing at least some of said equine sperm cells from said male species of said equine mammal comprises the steps of:

a. determining the sex characteristic of a plurality of said equine sperm cells; and

- b. sorting said equine sperm cells according to the determination of their sex characteristic,
and wherein said step of producing an equine offspring mammal comprises the step of producing an equine offspring mammal of the desired sex.

5 115 A method of practically producing an equine mammal as described in claim 111 wherein said step of establishing an equine insemination sample containing at least some of said equine sperm cells from said male species of said equine mammal comprises the steps of:

- 10 a. determining the sex characteristic of a plurality of said equine sperm cells; and
b. sorting said equine sperm cells according to the determination of their sex characteristic,
and wherein said step of producing an equine offspring mammal comprises the step of producing an equine offspring mammal of the desired sex.

15 116 A method of practically producing an equine mammal as described in claim 115 wherein said steps of determining the sex characteristic of a plurality of said equine sperm cells and sorting said equine sperm cells according to the determination of their sex characteristic comprise the steps of:

- 20 a. staining said equine sperm cells;
b. sorting according to said sex of said equine sperm cells through the use of high speed flow cytometry; and
c. concentrating said sorted equine sperm cells.

25 117 A method of practically producing an equine mammal as described in claim 116 wherein said step of sorting according to said sex of said equine sperm cells through the use of high speed flow cytometry comprises the step of collecting live sperm of the desired sex at the rate of at least nine hundred live sperm per second.

118 A method of practically producing an equine mammal as described in claim 116 wherein said step of sorting according to said sex of said equine sperm cells through

the use of high speed flow cytometry comprises the step of operating a high speed cell sorter at a pressure of at least about fifty pounds per square inch.

119 A method of practically producing an equine mammal as described in claim 116

wherein said step of sorting according to said sex of said equine sperm cells through the use of high speed flow cytometry comprises the step of collecting equine sperm cells having the desired sex characteristic in a skim milk solution.

120 A method of practically producing an equine mammal as described in claim 109 and further comprising the step of administering an equine pituitary extract to said female equine mammal to enhance the probability at which said step of fertilizing at least one equine egg within said female species of said equine mammal occurs.

121 A method of practically producing an equine mammal as described in claim 109 wherein said step of establishing an equine insemination sample containing at least some of said equine sperm cells from said male species of said equine mammal comprises the step of establishing an equine insemination sample having a volume selected from the group: 0.2 ml, or 1ml.

122 A method of practically producing an equine mammal as described in claim 110 wherein said step of establishing an equine insemination sample containing at least some of said equine sperm cells from said male species of said equine mammal comprises the step of establishing an equine insemination sample having a volume selected from the group: 0.2 ml, or 1ml.

123 A method of practically producing an equine mammal as described in claim 92 wherein said step of non-surgically inserting at least a portion of said equine insemination sample in said female species of said equine mammal comprises the step of inserting at least a portion of said equine insemination sample deep within at least one of said uterine horns of said female species near the tip of said uterine horn.

124 A method of practically producing an equine mammal as described in claim 109 wherein said step of non-surgically inserting at least a portion of said equine insemination sample in said female species of said equine mammal comprises the step of inserting at least a portion of said equine insemination sample deep within at least one of said uterine horns of said female species near the tip of said uterine horn.

125 A method of practically producing an equine mammal as described in claim 111 wherein said step of non-surgically inserting at least a portion of said equine insemination sample in said female species of said equine mammal comprises the step of inserting at least a portion of said equine insemination sample deep within at least one of said uterine horns of said female species near the tip of said uterine horn.

126 A method of practically producing an equine mammal as described in claim 110 wherein said step of non-surgically inserting at least a portion of said equine insemination sample in said female species of said equine mammal comprises the step of inserting at least a portion of said equine insemination sample deep within at least one of said uterine horns of said female species near the tip of said uterine horn.

127 A method of practically producing an equine mammal as described in claim 125 and further comprising the step of manipulating the ovulation of said female equine mammal prior to accomplishing the step of artificially inseminating said female species of said equine mammal.

128 A method of practically producing an equine mammal as described in claim 126 and further comprising the step of manipulating the ovulation of said female equine mammal prior to accomplishing the step of artificially inseminating said female species of said equine mammal.

129 A method of practically producing an equine mammal as described in claim 127 wherein said step of manipulating the ovulation of said female equine mammal prior to accomplishing the step of artificially inseminating said female species of said

equine mammal comprises the step of administering a gonadotropin releasing hormone to said female equine mammal.

5 130 A method of practically producing an equine mammal as described in claim 128 wherein said step of manipulating the ovulation of said female equine mammal prior to accomplishing the step of artificially inseminating said female species of said equine mammal comprises the step of administering a gonadotropin releasing hormone to said female equine mammal.

10 131 A method of practically producing an equine mammal as described in claim 129 wherein said step of artificially inseminating said female species of said equine mammal is accomplished at a time selected from the group consisting of: about thirty four hours after said step of administering said gonadotropin releasing hormone to said female equine mammal, about forty hours after said step of administering said gonadotropin releasing hormone to said female equine mammal, and between about thirty four hours to about forty hours after said step of administering said gonadotropin releasing hormone to said female equine mammal.

15 132 A method of practically producing an equine mammal as described in claim 130 wherein said step of artificially inseminating said female species of said equine mammal is accomplished at a time selected from the group consisting of: about thirty four hours after said step of administering said gonadotropin releasing hormone to said female equine mammal, about forty hours after said step of administering said gonadotropin releasing hormone to said female equine mammal, and between about thirty four hours to about forty hours after said step of administering said gonadotropin releasing hormone to said female equine mammal.

20 133 A method of practically producing an equine mammal as described in claim 129 wherein said step of inserting at least a portion of said equine insemination sample deep within at least one of said uterine horns of said female species near the tip of said uterine horn comprises the steps of:

25

- a. establishing a flexible probe having a sperm container;
- b. placing said flexible probe in the vagina of said female equine mammal;
- c. manipulating said flexible probe into said uterus of said female equine mammal;
- d. slowly guiding said flexible probe into a uterine horn of said female equine mammal; and
- e. gently manipulating said flexible probe per rectum as it is guided deep within said uterine horn of said female equine mammal to a location deep within said uterine horn of said female species near the tip of said uterine horn.

10 134 hod of practically producing an equine mammal as described in claim 130 wherein said step of inserting at least a portion of said equine insemination sample deep within at least one of said uterine horns of said female species near the tip of said uterine horn comprises the steps of:

- a. establishing a flexible probe having a sperm container;
- b. placing said flexible probe in the vagina of said female equine mammal;
- c. manipulating said flexible probe into said uterus of said female equine mammal;
- d. slowly guiding said flexible probe into a uterine horn of said female equine mammal; and
- e. gently manipulating said flexible probe per rectum as it is guided deep within said uterine horn of said female equine mammal to a location deep within said uterine horn of said female species near the tip of said uterine horn.

135 An animal produced through use of a method as described in any of claims 109, 113, 114, 117, 119, 120, 121, 122, 123, 127, or 133 .

1/2




Fig. 1

2/2




Fig. 2

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/17165

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : Please See Extra Sheet.

US CL : 424/811

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/811; 435/7.21; 435/806; 530/852; 800/8

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Please See Extra Sheet.

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Please See Extra Sheet.

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	MCKINNON, James L. et al. Equine Reproduction. Philadelphia: Lea & Febiger. 1993, pages 291, 345-347, 738-754, 782-789, 793-796.	1, 9, 10, 14-21, 24, 25, 29, 30, 35, 38, 39, 40, 42, 44-47, 52-59, 61, 62, 64, 68-135
---		---
Y		2-8, 11-13, 22, 31-34, 36, 37, 39, 43, 48-51, 60, 63, 65-67
---		---
A		27, 28, 41

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
B earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

13 SEPTEMBER 1999

Date of mailing of the international search report

27 OCT 1999

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 308-4242

Authorized officer

KRIS PELHAM MAYO

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/17165 ..

C (Continuation). DOCUMENTS--CONSIDERED--TO--BE--RELEVANT--		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X, P	VAZQUEZ, J.J. et al. 'Nonsurgical Uterotubal Insemination in the Mare.' In: Proceedings of the 44th Annual Convention of the American Association of Equine Practitioners. Lexington, Kentucky: AAEP, 1998, pages 68-69.	1, 3-21, 24, 25, 27-44, 52-55, 58, 59, 70-135
X	BRINSKO, Steven P. et al. Artificial Insemination and Preservation of Semen. The Veterinary Clinics of North America, Equine Practice. April 1992, Vol.8, No.1, pages 207-211.	1, 9, 10, 14-21, 24, 25, 29, 30, 35, 38-40, 42, 44-47, 52-59, 61, 62, 64, 68, 69, 70-135
---		---
A		3-8, 11, 27, 28, 31-34, 36, 37, 41, 43
X	MEINERT, C. et al. Advancing the time of ovulation in the mare with a short-term implant releasing the GnRH analogue deslorelin. Equine Veterinary Journal. January 1993, Vol.25, No.1, pages 65-68.	21,22,25,40,83-88,120,121,127-135
X	GRAHAM, James. Analysis of Stallion Semen and its Relation to Fertility. The Veterinary Clinics of North America, Equine Practice. April 1996, Vol.12, No. 1, pages 127-130.	2-14,31-39,43-55,58-69,71-74,76,78,101-108,114-119,135
X	RATH, D. et al. Production of piglets preselected for sex following in vitro fertilization with X and Y chromosome-bearing spermatozoa sorted by flow cytometry. Theriogenology. 1997, Vol.47, pages 795-800.	2-14,31-39, 43-55,58-69,71-74,76,78,101,108, 114-119,135
X	US 5,439,362 A (G.F. SPAULDING) 08 August 1995, abstract, column 3, lines 24-34, column 4, lines 28-36, column 7-10, Example I, claim-3.	2-14,31-39,43-55,58-69,71-74,76,78,101-108,114-119,135
X	US 4,999,283 A (P.M. ZAVOS et al.) 12 March 1991, column 1, lines 9-19.	2-14,31-39,43-55,58-69,71-74,76,78,101-108,114-119,135

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/17165 --

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 4,339,434 A (R.J. ERICSSON) 13 July 1982, column 1, lines 26-64, column 2, lines 5-29.	2-14,31-39,43-55,58-69,71-74,76,78,101-108,114-119,135
X	WELCH, G.R. Flow Cytometric Sperm Sorting and PCR to Confirm Separation of X- and Y- Chromosome Bearing Sperm. Animal Biotechnology. 1995, Vol.6, No.2, pages 131-139, especially pages 131, 133.	2-14,31-39, 43-55,58-69,71-74, 76,78,101-108,114-119,135
X	PICKETT, B.W. et al. Recent developments in artificial insemination in horses. Livestock Production Science. 1994, Vol. 40, pages 31-36, especially pages 32, 34, and 35.	3-14,27-37, 41-47,53-55,58,59, 61,62,64, 68,69,74-78,92,94-135
X	JOHNSON, L.A. et al. Modification of a Laser-Based Flow Cytometer for High-Resolution DNA Analysis of Mammalian Spermatozoa. Cytometry. May 1986, Vol.7 No.3, pages 268-273.	2-14,31-39, 43-55,58-69,71-74, 76,78,101-108
X	PINKEL, D. et al. Flow Cytometric Determination of the Proportions of X- and Y-Chromosome-Bearing Sperm in Samples of Purportedly Separated Bull Sperm. Journal of Animal Science. 1985, Vol.60, No.5, pages 1303-1307.	2-14,31-39, 43-55,58-69,71-74, 76,78,101-108

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/17165

A. CLASSIFICATION OF SUBJECT MATTER:

IPC (6):

A61K 39/00, 39/395

B. FIELDS SEARCHED

Documentation other than minimum documentation that are included in the fields searched:

McKinnon and Voss; AAEP Proceedings, 1998; The Veterinary Clinic of North America 8:1; The Veterinary Clinics of North America 12:1

B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

STN, AGRICOLA, CAPLUS, MEDLINE, BIOSIS, CABA, USPATFULL, WEST

SEARCH TERMS: horse, equine, sperm, sperm sorting, sex selection, gender selection, artificial insemination, technique, uterine horn, tip, UTJ, uterotubal junction, flow cytometry, deslorelin